

Studies on the Johnstonianidae (Acari, Parasitengona)

IRWIN M. NEWELL¹

THE SUBFAMILY Johnstonianinae Thor 1935 (Trombidiidae) was established to include *Johnstoniana* George 1909 and related genera. The only comprehensive treatment of the group was published by Thor and Willmann (1947) who included eight species in the genera *Centrotrombidium* Kramer 1896, *Johnstoniana*, *Diplothrombium* Berlese 1910, *Myrmicotrombium* Womersley 1934, and *Hirstithrombium* Oudemans 1928.

The present study was undertaken with a view to placing the systematics of the Johnstonianinae on a sound morphological basis, and to determine the relationship of the group to other terrestrial Parasitengona. It soon became apparent that the mites included in the Johnstonianinae differed so significantly from other terrestrial Parasitengona that they could not logically be retained within the family Trombidiidae Murray 1877.² The differences are much greater than those which have been cited, for example, to establish Trombiculidae Ewing 1944 as a family separate from the Trombidiidae. The group is therefore raised to family rank with *Johnstoniana* George 1909 as the type genus.

Interest in the group was stimulated by numerous indications that this is perhaps the most primitive existing family, terrestrial or otherwise, within the Parasitengona. The indications of this are of ecological, behavioral, and morphological nature.

Ecologically the Johnstonianidae are terrestrial in larval, nymphal and adult stages, but are rarely found where there is not an ample supply of water nearby; in fact many of them could be aptly termed subaquatic. Thus, ecologically they are in a position from which they (or their antecedents) could evolve in two directions—either toward strictly terrestrial forms such as the Trombidiidae, or toward the subterrestrial aquatic mites such as the Limnocharidae, Thyasidae, etc., and thence to the more strictly aquatic Parasitengona. From an evolutionary standpoint, it might be hypothesized that the Parasitengona parasitic on insects evolved before those on terrestrial vertebrates, because insects appeared first in the fossil record. This assumes that there were appropriate predatory Acari present when the hosts appeared, but since the earliest fossil record of the Trombidiiformes (*Protacarus* Hirst, Devonian) is at least contemporaneous with that of the earliest known Insecta, this is not an impossible assumption. The next major group of hosts to become available to the Parasitengona were the Amphibia, and the seemingly close structural similarities between *Hannemania* species (which largely parasitize Amphibia) and the Johnstonianidae, which parasitize insects living in moist habitats, is suggestive that these are phylogenetically very close. This close relationship could have arisen in more than one way—either by the direct descent of the Amphibia-parasites from the early Johnstonianidae, or by simultaneous radiation of the Amphibia-parasites and the Johnstonianidae from a common ancestral mite or group of mites which appeared at the same time as or somewhat later than the Amphibia. The extent to which the Parasitengona have become adapted to the Insecta

¹ University of California, Riverside, California. Manuscript received April 17, 1956.

² While many derivatives of the name *Trombidium* Fabricius 1775 were published prior to 1876 (see Oudemans 1937: 1349–1361), the first unequivocal use of the name Trombidiidae appears to be that of Murray (1876) in his *Economic Entomology*. Accordingly the writer ascribes the family name to Murray 1876 rather than to Leach 1815, who employed the name Trombides, and not Trombidiidae.

would suggest that the only thing that might possibly have prevented the parallel emergence of the latter two groups would have been the unavailability of a suitable prototype mite at the time the insects were first becoming established. The groups with larvae parasitic on higher vertebrates appear to be more modified than the Johnstonianidae and *Hannemania* (although not more so than the Trombidiidae or Erythraeidae which parasitize insects).

So far as their habits are concerned, the larvae of the Johnstonianidae are all parasitic, but the relationship between larva and host is very loose. Among the Parasitengona there are two extremes in the relationship of larva to host—in some species the larva is little more than a predator, preying upon a number of host individuals during its development, and detaching readily when disturbed. At the other extreme we find larvae which, once firmly attached, remain with the host until they complete their larval development almost regardless of what stimuli may intervene. A simple test of the degree of host fixity is to drop the host with the attached larvae into a vial of 60 per cent alcohol. Forms in which host fixity is highly developed will usually remain attached, unless they had not become firmly attached to the host or unless they were in the process of detaching themselves at the time of capture. Forms in the other group detach readily regardless of how long they have been on the host, and for this reason they are called "self-detaching larvae." All Johnstonianidae which the author has studied in the living state have larvae of the self-detaching type.

Morphologically they appear to be more generalized than other Parasitengona. For example, there are two pairs of sensilla on the scutum in most species, such as are found in the Erythraeidae and most Smarididae, but with obvious trends toward modification or loss of the anterior pair as in the Trombidiidae and Trombiculidae. The simplicity of the body setae and those of most leg segments

would appear to be a primitive characteristic contrasting with the more ornate setae found in other families. The possession of a large complement of supracoxal, vestigial, and rostral setae would also seem to be primary in the Parasitengona, secondary reductions in these being most notable in the Trombidiidae and Trombiculidae. It should be pointed out, however, that notable differences in these occur within the limits of the family. The possession of well-developed paragenital sclerites also seems to be a primitive characteristic. These are probably found in all genera of terrestrial Parasitengona, although in many they lack setae and are nearly invisible. The generalized structure of the palpal tarsus of the larva, and the presence of a single subterminal spiniform seta on the adult palpal tibia, appear in sharp contrast to the greatly reduced tarsus of the larvae of the Trombidiidae, and to the elaborately developed ctenidium of the adults of many genera outside of the Johnstonianidae. These are some of the principal morphological indications of the primitive nature of the group. The detailed morphology of these and other structures will be discussed below, along with others not mentioned here.

ACKNOWLEDGMENTS

The present study was initiated under a contract between the Office of Naval Research (Nonr-782(00)), and the University of Hawaii while the writer was on the staff of the Hawaii Agricultural Experiment Station. It was continued under a research grant (NSF-G1833) from the National Science Foundation to the University of California. Much of the work, including the field study program for 1953, has been supported by research grants from the University of California. Without the assistance of these foundations and institutions, the present work would have been most improbable.

Mari Riess of the University of California at Riverside made about 200 of the camera lucida drawings; while the remainder were

done by Helen Au Randall, formerly of the University of Hawaii. In these figures, the scales provided are marked in 10 μ units, except for the few marked in 100 μ units. The latter are drawn with a double base line, and also have the total length indicated on the scale.

This is not intended to be an exhaustive review of the species assignable to the family. While such might be desirable, as a matter of fact existing descriptions are so inadequate that a thorough review of the group is impossible until all species have been redescribed. Only where it was felt that a specific contribution to knowledge could be made on the basis of the descriptive material available has mention been made of previously named forms. For example, the genus *Polydora* Methlagl is shown to belong in the family, at least provisionally, and a brief mention of *Crossothrombium* Womersley, omitted from recent catalogues, is included to bring it to the attention of other workers who might also overlook it.

MORPHOLOGY

In describing these and other mites, the writer has found it useful to develop a system of notation of the position of specialized setae or other features on the segments of the legs and palpi. It is frequently necessary to indicate the precise position of these, but at present there is no brief, convenient method for doing so. The system will be illustrated by an example. The position of the seta or other structure is stated with reference to the proximal and distal ends of the segment. For instance, the statement "tarsus II with a spikelike famulus at 0.61 pd " means that famulus₂ is located at a point 0.61 of the length of the tarsus from the proximal end, and on the posterodorsal aspect of the segment. The terms dorsal, ventral, anterior, and posterior, are utilized to express position with relationship to the longitudinal axis of the legs. The terms lateral and medial are not

employed, since the meaning of these terms is reversed on legs I and II as compared with III and IV. The reader should visualize the legs as protruding from the sides of the body at right angles to the main axis of the body. For the sake of morphological uniformity, the same orientation should be visualized in the case of the pedipalps despite the seeming incongruity of the pedipalps of a mite projecting at right angles to the median axis of the body. The terms preaxial and postaxial are in some respects preferable to anterior and posterior, in referring to the segments of appendages, but the writer prefers to use anterior and posterior because the symbols a and p are immediately understandable, avoiding the necessity of using either two-letter symbols, or others which are less readily comprehensible. In the following descriptive accounts, the symbol a following a decimal in the designation of the position of a particular seta or structure means anterior, p = posterior, d = dorsal, v = ventral, ad = anterodorsal, pd = posterodorsal, av = anteroventral, and pv = posteroventral. This method of notation greatly simplifies the problem of precise location of setae, and also makes it possible to analyze variation in the positions of specific setae. By the use of special graphical methods it is possible to directly determine the positions of the specialized setae with ease and rapidity, without the necessity of making measurements and converting these to decimals.

In naming the specialized setae of the appendages, the writer has adopted the system first propounded by Grandjean, beginning in 1935. The only modification was the addition of a system of notations for the several types of solenidia found on the legs. The Grandjean system of setal nomenclature is based primarily upon the form and physical properties of the setae, rather than upon their position. The solenidia are thin-walled setae showing spiral or annular internal structure in most types, although some solenidia do not show any internal structure. Typically the

solenidia taper very little throughout their length, and the distal end is usually rounded. Ornamentation is almost invariably wanting, and the cuticle is optically inert in polarized light. The eupathidia are similar to the solenidia in certain respects, but there is usually no internal structure (an exception is illustrated in Fig. 238). External ornamentation is more likely to be found than in the solenidia, in the form of short barbs (Figs. 215, 216, 217) and the cuticle is nearly always partially anisotropic. Birefringence is most noticeable in the basal portion. The walls of the eupathidia are typically of uniform thickness throughout—slightly thicker than the solenidia and thinner, relatively, than the normal setae. Notable transformations in structure of the eupathidia are found in certain cases, as in those on the tarsus of the palp. For example, in the genera *Centrotrombidium*, *Johnstoniana*, and *Diplothrombium*, typical eupathidia are found on the palpal tarsus of the adult (Figs. 59, 83, 159), while in the larva the apparent ontogenetic forerunners of these are heavy, pectinate, sometimes elaborately formed setae, with little or no trace of a central canal (Figs. 29, 92, 132).

The other types of setae provide no particular difficulties in interpretation and will be discussed below with reference to their characteristics and distribution in the Johnstonianidae.

Scutum

A number of interesting and illuminating trends can be seen in the structure and chaetotaxy of the scutum of the Johnstonianidae. The first is a reduction or a modification of the anterior pair of sensilla, leading to their eventual disappearance, or their transformation into setae which are not at all sensillar in form. While it might be argued that the trend is the reverse, that is, toward the development of the anterior pair of sensilla, this does not appear to be the most likely alternative. In *Johnstoniana* there is a well defined anterior pair of sensilla in both larva and adult

(Figs. 70, 90). Also, in the larva of *Lassenia scutellata* there is a well defined anterior pair of sensilla, borne on a small sclerite separated from the remainder of the scutum by a narrow band of striated membranous cuticle (Fig. 249). In the adults of *Lassenia spinifera*, these anterior setae (Fig. 232) are scarcely recognizable as sensilla because the alveoli are not to any sensible degree different from those of the other setae of the scutum. But there is little doubt that the two short simple setae on the spine of the adult scutum actually are the anterior sensilla. In the larva of *Lassenia laseni* the anterior sensilla are present, although the alveoli of these are not as well developed as the alveoli of the posterior pair of sensilla (Fig. 214). Nevertheless these are unquestionably the anterior sensilla. They are short and distinctly barbed, in marked contrast to the posterior pair which are long, slender, and smooth. In fact the anterior sensilla more closely resemble the other setae on the scutum than they do the posterior sensilla. In the adult of the species (Fig. 183) there is only one well defined pair of sensilla. At the same time there is usually a pair of stiff, rodlike setae near the anterior margin of the scutum, somewhat set apart from the rest of the setae. If one were to examine only the nymphs and adults of *Lassenia laseni* there would be considerable justification in stating that this species was characterized by the possession of a single pair of sensilla. Yet there is no doubt that the anterior two rodlike setae on the adult scutum are derived from the anterior sensilla of the larva, and are therefore homologous with them, regardless of their remarkable transformation in form. Thus within the one genus *Lassenia* we see a partial obscuring of the anterior pair of sensilla between the larval and adult stages.

Genera in other families of Parasitengona should be re-examined to determine those in which the anteromedian setae of the scutum are homologous with the anterior sensilla for it is evident from the Johnstonianidae that this can be expected. Such is almost certainly

the case in *Hannemania*, *Leeuwenhoekia*, and related genera in which the resemblance to some of the Johnstonianidae is augmented by the possession of an anteromedian spine on the scutum. If these are homologous with the anterior sensilla, then it is further likely that the single anteromedian seta found in many of the Trombiculidae is also homologous with the anterior sensilla, which have somehow or other become fused. Only in such genera as *Gateria*, *Walchia*, *Gabrliepia* and *Schöngastiella* do the anterior sensilla (or their paired or unpaired homologues) appear to be totally lacking. The recognition of the essential unity of these setae at the anterior end of the scutum will help greatly in the analysis of evolutionary trends within the Parasitengona. The modifications of the anterior sensilla in the Johnstonianidae provide the key.

In *Diplothrombium monoense*, we find that it is the adult rather than the larva in which the anterior sensilla are best developed (Figs. 110, 129). And, finally, in *Centrotrombidium* the anterior sensilla are completely absent in both larva and the adult. Some authors have referred to the anterior pair of setae on the scutum of *Centrotrombidium* as a second pair of sensilla, but a comparison of Figures 4 and 110 will show that this is not the case; rather the anterior pair of setae in *Centrotrombidium* are undoubtedly the homologues of the setae which, in *Diplothrombium*, lie between the two pairs of sensilla; the anterior sensilla are totally lacking.

Another interesting variant is found in the degree of duplication of the setae of the scutum other than the sensilla. In *Centrotrombidium* and *Diplothrombium*, the setae anterior to the posterior sensilla undergo little or no reproduction in the transformation from the larva to adult, except perhaps in *D. micidium*. The larva of this species is as yet unknown (unless it should prove to be the larva described here as *D. cascadenae*) so it is not possible to state positively whether or not there has been a duplication of these setae in

this particular species. If we assume that the larva does have a single pair of setae between the anterior and posterior pairs of sensilla, which is probably the case, then there is a moderate degree of duplication of these setae (Fig. 160).

In *Diplothrombium monoense* there is a moderate degree of duplication of these setae as a rule, for there is usually a second pair of setae near the margin of the plate between the anterior and posterior pairs of sensilla (Fig. 110). Considering the situation in *D. micidium*, it is probable that these can be regarded as a duplication of the setae in question. The situation in *Johnstoniana* and *Lassenia* is somewhat more complex, for there are usually five to nine or more setae which we could regard as the ontogenic descendants of the intersensillar setae of the larva.

In addition to the chaetotactic features described above, there is a further variant in the structure of the scutum. In both *Centrotrombidium* and *Diplothrombium*, there is a noticeable tendency toward the formation of a posterior stalk with a feebly sclerotized lateral expansion in the adult (Figs. 4, 63, 110, 160). No corresponding development is found in the scutum of either *Lassenia* or *Johnstoniana*. Associated with the tendency toward the differentiation of the posterior lobe of the scutum, is the degree of development of the lateral portion of the scutum outside the crista. The broadest scutum is found in *Johnstoniana*, a somewhat narrower plate in *Lassenia*, and the narrowest scuta are found in *Diplothrombium* and *Centrotrombidium*.

There is also a noticeable difference in the crista metopica, for this structure is well developed in the larva of *Johnstoniana* and *Diplothrombium*, very feebly developed in the posterior part of the scutum of *Centrotrombidium*, and essentially absent from the larval scutum of *Lassenia*. A unique variant in the form of the posterior sensilla is found in *Centrotrombidium*, in which the distal portion of the shaft is greatly expanded and pyriform to spherical in form.

Corneae

The eyes of the Johnstonianidae show little variation. There are typically two well-developed corneae borne on a small ocular plate on either side of the scutum. While these frequently protrude prominently, they are not so well developed that they could be described as stalked, except in *Johnstoniana*, in which the ocular plates are definitely cylindrical and raised well above the surface of the propodosoma (Fig. 86). About the only variant in the form of the corneae is found in *Centrotrombidium*. In this genus only the anterior cornea is developed in the adults of the species known to the writer; the posterior cornea is represented by a prominent but nonhyaline protuberance (Fig. 53). In none of the forms seen by the writer are there any setae on the ocular plates, either in the larva or in the adult. According to Womersley, the eyes are absent in *Crossothrombium*.

Dorsal Setae of Hysterosoma

There are two outstanding features of the dorsal chaetotaxy of the Johnstonianidae. The first of these is the insertion of the dorsal setae of the larvae in individual sclerites which show no sign of fusion in any of the known forms. The second feature is the rather poor development of setal ornamentation, especially in the adults. As in the larva the setae of the adults are also borne on individual sclerites, although these are relatively smaller than in the larva. In all species studied, the dorsal setae in the adult are completely smooth, and are frequently stiff, short, and rodlike (Fig. 183). In the larva the setae are always arranged in four or five well defined rows (Fig. 131). The dorsal setigerous sclerites in the larva are always flat or weakly arched, but in the adult they are characteristically elevated, forming either low hemispheres or short cylindrical protuberances above the general surface of the cuticle (Fig. 64).

Dorsal Propodosomal Setae

In none of the known larvae are there any

setae outside the scutum. In the adult however, there is a tendency toward the development of setae lateral to the scutum and even lateral to the ocular plate. In *Johnstoniana latiscuta* (Fig. 86) there are no setae whatever lateral to the scutum. In *Lassenia lasseni* there is a single pair of setae, or occasionally more, between the scutum and the ocular plate, but the great majority of the propodosoma outside of the scutum is bare (Fig. 183). The situation in *Lassenia spinifera* is somewhat different, for in this species there are quite a number of setae surrounding the ocular plate although these are by no means as abundant as on the dorsum of the hysterosoma; and moreover there is an appreciable difference in form between the dorsal setae of the propodosoma and hysterosoma (Fig. 232). In *Centrotrombidium* and *Diplothrombium*, there are generally several setae between the scutum and ocular plates, although again the density of setae here is in no way comparable with that on the dorsum of the hysterosoma.

Coxal Chaetotaxy of Larva

The situation with respect to this character is not entirely clear owing to the small number of species known in each of the genera. In each of the four species of *Centrotrombidium*, *Diplothrombium*, and *Johnstoniana* which have been adequately described there are two setae on coxa I and one seta on each of coxae III and IV (2-1-1). Only two species are known in the genus *Lassenia*, the coxae of *L. lasseni* having the setal formula 2-2-3, and *L. scutellata* having the setal formula 2-1-2. Thus it is possible that in the genus *Lassenia* the number of coxal setae in the larvae is only a specific character, while in the other three genera it may prove to be generically constant. The supracoxal setae of the legs are discussed below along with the other specialized setae of the appendages.

Pars Medialis Coxae

This is the name applied to the small extension of the medial angle of coxa I of most

adult terrestrial Parasitengona (Figs. 54, 82, 154, 192). In many cases, especially in the adult, it is difficult to tell whether this is a part of coxa I or II, or whether it is a distinct sclerotized area independent of the coxae. However, in those species in which coxae I and II are rather distinctly separated, and in which a distinct pars medialis coxae is developed, it is clear that this is an extension of the coxa of leg I. In those cases in which the relationship is not so clear, it is probably safe to assume that the pars medialis coxae is morphologically a part of coxa I rather than II.

The number of setae present on the pars medialis coxae is quite variable even within the limits of a given species. At best, variations in this structure appear to be of a specific rather than a generic nature. An interesting variant is found in the genus *Lassenia*, in which *L. lasseni* has a well-defined pars medialis coxae in the adult (Fig. 192) whereas in *L. spinifera* (Fig. 234) the pars is entirely absent. In the latter species there is a general reduction in the degree of sclerotization of the coxae for many of the setae which normally would be included within coxae II actually lie in the membranous cuticle immediately behind the posterior margin of these (Fig. 234). It is interesting to note that when there is a difference in the form of the setae of coxae I and the intercoxal area such as exists in *Diplothrombium micidium* (Fig. 154) and *Lassenia lasseni* (Fig. 192) the setae of the pars medialis coxae are more similar to those of the intercoxal area than they are to those of the coxae themselves.

Coxal Ring

In the Parasitengona, the coxae have become largely incorporated into the ventrolateral body wall as a series of well-defined plates. However, if one examines the distal portion of the typical coxa of an adult it is found that this comprises a cylinder of extremely short length, and it is this peripheral portion which the author refers to as the coxal

ring. The distal portion of the coxa is ringlike in the adults of most species of terrestrial Parasitengona, except for leg I, in which the coxal ring is generally, if not always, incomplete dorsally (Fig. 107b). This is the situation in the adults of all genera except *Lassenia*, in which the two known species have both coxal rings I and II incomplete dorsally in the adult (Fig. 232). In all species of all genera of the Johnstonianidae, coxal rings III and IV are complete dorsally (Fig. 107a).

Urpore

The urpore is present in the larvae of all known species of Johnstonianidae and is well developed. There are no variants of known generic significance.

Lassenia Organ

This is a term which the author has applied to a structure of unknown nature and function immediately anterior to coxa III of larvae and adults of certain of the Johnstonianidae. It is especially well developed in *Lassenia lasseni* and *Lassenia scutellata* (Figs. 199, 209, 221, 251). A search of specimens of *Johnstoniana* and *Centrotrombidium* has failed to reveal a corresponding organ in these genera. At least a rudiment of this organ is found in larvae of *Diplothrombium monoense*, and *D. cascadeense*, but not as yet in the adults of these species. A systematic search of other Parasitengona might reveal other groups in which this organ is found. It is most likely a gland of unknown function for in a number of specimens, both larva and adult, a duct leads from the body surface into the incompletely hydrolyzed remains of a glandlike mass of cells. Beyond this its nature is unknown.

Genital and Paragenital Sclerites of Adult

The Johnstonianidae are somewhat unusual among the Parasitengona in that the genital and paragenital sclerites are almost equally developed. Thus the genital opening is surrounded by four sclerites of very nearly equal size, and bearing roughly equal numbers of

setae. In fact, in *Diplothrombium micidium* the setae on the paragenital sclerites actually outnumber those on the genital sclerites proper. In the other terrestrial Parasitengona, the paragenital sclerites bear markedly fewer setae than the genital sclerites, if they bear any at all. There are invariably three pairs of genital acetabula in both the male and female. The penis of the male is relatively very small. A unique variant in the genital area is found in *Lassenia* in which there is a well developed pregenital tubercle. In *Lassenia lassemi* this makes its first appearance in the protonymph and increases in size in the deutonymph and adult. In *L. lassemi* it is a simple hemispherical tubercle, while in *L. spinifera* it is an elongate peduncle. In both cases it is heavily sclerotized and pigmented in the mature adult. Not even a rudiment of this pregenital tubercle has been found in the larvae, nymphs or adults of any of the other genera.

Anal Area

In all genera but *Lassenia* and *Polydiscia* the anal anlage of the larva is a simple slit with no associated setae. In both *Lassenia lassemi* and *L. scutellata* the slit is guarded by a pair of anal sclerites bearing two pairs of simple, smooth setae. The anal sclerites of the adults of all but *Johnstoniana* are distinctively crescentic in form, bearing in most cases a single row of smooth, simple setae (Figs. 18, 105, 240). In *Lassenia spinifera* there are two rows of setae on the anal sclerites, whereas in at least one male of *Lassenia lassemi*, the two anal sclerites had only 0 and 1 setae respectively. Perhaps the most significant variant is found in *Johnstoniana latiscuta* in which there are no anal sclerites at all (Fig. 68). There is a poorly defined anal area bearing a number of setae and differing from the surrounding cuticle only in the absence of striae.

Rostral Setae

The situation with regard to these setae cannot be finally resolved until further studies are made in other families. However, if we

take *Lassenia lassemi* as a starting point, we find in the larva of that species three pairs of setae on the gnathosoma, all of these on the rostrum (Fig. 229). These will be called the proto-, deuto-, and tritorstral setae, and the presence of these three pairs is a condition characteristic of the larvae of many genera of the terrestrial Parasitengona. The protorostral setae are generally located dorsally or dorso-laterally on the tip of the rostrum, not only in this species but in the terrestrial Parasitengona in general. (These are the so-called "galeal setae" of specialists in the Trombiculidae. The writer prefers the term "protorostral setae" because it is simpler to give these three topographically related setae parallel names than to coin separate ones for each.) In *L. lassemi*, the deutorostral setae are under the margin of the velum in both the larva and the adult (Figs. 188, 229), while the tritorstral setae lie directly behind them. In *Centrotrombidium distans*, *Diplothrombium* and *Johnstoniana* there are only two pairs of setae in the larva (Figs. 45, 170, and 88). The distal pair of setae obviously are the protorostrals while the basal pair are either deutorostrals or tritorstrals. In larvae of *Centrotrombidium distans* (Fig. 45) a pair of very minute structures of uncertain nature is found along the margin of the velum in approximately the position occupied by the deutorostral setae of *Lassenia lassemi*. Apparently homologous structures are found in the adult as well (Figs. 2, 17). These may represent highly modified setae, or they may simply be points of insertion of muscles associated with the velum. Homologous structures have been found also in adults of *Diplothrombium monoense*, but neither in the adults nor the larvae of *Johnstoniana latiscuta*. From their position alone these would appear to be homologous with the deutorostral setae of *Lassenia* so that the series *Lassenia*—(*Centrotrombidium*, *Diplothrombium*)—*Johnstoniana* represents a progressive diminution in the size and importance of the deutorostral setae. This conclusion should be verified with studies in related genera and

families, but at present this appears to be the situation in the Johnstonianidae. In none of the larvae of the Johnstonianidae studied so far are there any setae behind the tritrostrals corresponding to the posterorostrals of some genera.

In the adults, of course, the base of the gnathosoma is generally covered with a considerably greater number of setae. An interesting exception is found in *Lassenia spinifera* (Fig. 248), in which only one pair of setae is added behind the tritrostrals; in *L. lasseni* (Fig. 180) many more setae are found here. In adults of *Centrotrombidium* (Fig. 17) the number of setae is apparently never great, but these are small species. The largest number of setae found behind the tritrostrals is in *Diplothrombium* and *Johnstoniana* (Figs. 125, 66).

It is difficult to say whether or not the setae found behind the tritrostrals are to be considered the ontogenic descendants of the tritrostral setae, but this does not appear to be the case, for the tritrostrals often retain a fairly characteristic appearance, differing rather markedly from the setae behind them (Figs. 23, 106). We would therefore have to consider these setae as having arisen *de novo* in the postlarval instars.

Velum

This structure appears to show few variants of any significance in the family. It is a simple structure of moderate size in all species studied by the writer.

Posterolateral Arms of Gnathosoma

Only moderate trends are noticeable within the family, with *Johnstoniana* and *Diplothrombium* having essentially no lateral arms on the posterior margin of the gnathosoma and with *Centrotrombidium* having only moderately developed arms here. The maximum development of these structures is found in *Lassenia*. Within this genus there is quite a bit of difference between the two known species, with *L.*

spinifera having relatively longer posterolateral arms than *L. lasseni*.

Chelicerae

The form of the chelicerae in the Johnstonianidae is fairly uniform, and is best seen in Figures 74 and 85. The tarsus or digitus mobilis is scythe-shaped with the dorsal margin ranging from nearly smooth to serrate (Fig. 7) to dentate (Fig. 247). The digitus fixus is membranous and shows no particular variants of generic importance. Perhaps the most significant variant in the form of the chelicerae is found in the tendency toward a downward flexure in the chelicerae of adults of the genus *Lassenia*. In *Lassenia lasseni* this flexure is only slightly developed, but in *L. spinifera* it is very pronounced (Figs. 200, 245). One very interesting bilateral anomaly was found in a single female of *Centrotrombidium distans* (Fig. 24) in which a typical seta and alveolus were found dorsally at the base of the digitus fixus. No other individuals of this species were found to have such a seta here, and certainly none of the other Johnstonianidae studied had a seta in this position, or anywhere on the chelicera. Such a seta is found at this point in many genera of *Eleutherengona*, however, and also in many of the Parasitiformes. The appearance of this seta in this anomalous individual not only suggests that the Johnstonianidae were derived from a group which did have a seta on the chelicera, but also raises the question why such a structure, apparently so irrevocably lost in the course of evolution of the group, should suddenly reappear as a bilateral variant in a single female of *Centrotrombidium distans*. Further details of this will be found under the description of the species.

Segmentation and General Form of the Palp

With a single exception the palpi of the Johnstonianidae are provided with five free segments. In larvae of *Lassenia lasseni* (Figs. 210, 213) there is a tendency toward the fusion of the femur and patella on the dorsal side;

this is not seen however in *L. scutellata* so that the character does not seem to have generic significance. In the adults the palpi are always distinctly five-segmented. The general form of the palpi in the adult is basically the same in all genera, slightly curved but essentially linear in dorsal view. The same is true of the larvae of all of the genera except *Lassenia*, in which the palpi are geniculate, owing to the expansion of the posterior or posterolateral aspect of the femur. Associated with this expansion of the femur, the trochanter is reduced to a narrow ring. The geniculate palpi of *Lassenia* larvae are suggestive of similar palpi which show up at other points within the terrestrial Parasitengona, and in these cases too it is interesting to note that the geniculate form is lost in the transition from larval to nymphal stages.

Fenestration of Trochanter of Palp

This is a characteristic which presently appears to be confined to the family Johnstonianidae. In the larva and adult of *Diplothrombium* and *Centrotrombidium* (Figs. 33, 61, 153, 179) there is a discrete oval window on the anterior (medial) aspect of the trochanter of the palp. This appears to be a portion of the cuticle which is considerably thinner and more transparent than the cuticle of the remainder of the trochanter. In *Johnstoniana* there is no trace of a fenestra. In *Lassenia* the situation is somewhat variable. In both of the known species the larva has a trochanter of very short length, probably associated with the geniculate form of the palp. In this there is no suggestion of a fenestra. In the adult of *L. lasseni* (Fig. 201) there is a well-developed fenestra, but the anterior margin is not closed off as it is in *Diplothrombium* and *Centrotrombidium*. Thus the trochanter has the appearance of being deeply incised on the anterior aspect. In *L. spinifera* (Fig. 239) the same condition is found, except that here the concavity is invaded by a distinct, although feeble extension of the cuticle of the femur.

The Palpal Tibia

The form of the tibia in the Johnstonianidae known to date is quite constant in the larva. In all species there is a heavy terminal seta which is unidentate in *Lassenia scutellata* and *Centrotrombidium distans*, but bidentate in the other four species for which larvae have been adequately described. The terminal setae have been given a number of names by workers in various divisions of the Parasitengona including "tibial claw, tibial spur," etc. While the name is not of paramount importance, it is well to keep in mind that this is not a claw, but nothing more nor less than a seta of unusual thickness, and it is preferable to refer to it as the terminal seta of the tibia. The term "claw" is a malapropism. In the larvae of all known species there are three normal setae behind the heavy spiniform terminal seta, and these may be either smooth or faintly pectinate.

In the adults of all known species, the terminal seta of the tibia is invariably unidentate. Moreover there is always a single subterminal spiniform seta which may be very close to the terminal seta as in *Lassenia* and *Diplothrombium micidium*, or may be removed from it by a distance greater than the diameter of the alveolus of the terminal seta as in *Johnstoniana* and *Diplothrombium monoense*. In *J. latiscuta* there is a heavy spine near the base of the subterminal spiniform seta. In some of the Trombidiidae there is also a *ctenidium* composed of several stout setae arranged in a regular series, but this structure is not present in any of the known Johnstonianidae.

Segmentation of the Legs of the Larvae

In both species of the genus *Lassenia* the femur is completely undivided so that there are only five free segments beyond the coxa. In *Johnstoniana latiscuta*, there is a distinct synarthrodial membrane on the ventral surface of the femur, but the dorsal sclerotized portion of the cuticle is continuous across the entire length of the femur (Figs. 87, 91, 93).

In both *Centrotrombidium* and *Diplothrombium* the basifemur and telofemur are completely separated by a flexible synarthrodial membrane. The femora of the adults of all species are divided.

Specialized Setae of the Appendages

Here it is well to digress a little in order to point out a difficult situation which exists in the nomenclature of the specialized setae of the appendages of the Parasitengona. In the United States the majority of the work in the terrestrial Parasitengona has been done in the Trombiculidae, by investigators whose major or even sole interest in the group was stimulated by the medical or general parasitological importance of these mites. In no case is there any published evidence that these workers have availed themselves of Grandjean's studies on the types of setae found on the appendages of mites in general including the Parasitengona. Grandjean's unexcelled contributions to the chaetotaxy of the Acari began in 1935 and have continued up to the present time. The major setal types were clearly elucidated and named prior to 1940. Grandjean's studies were climaxed in 1947 with the publication of his *Étude sur les Smarisidae* (1947) in which the broad outlines of the morphology of the various setal types on the appendages of the Parasitengona were laid down. This was one of the classics of modern acarology and it is incomprehensible that certain workers should have proposed a "standardized terminology" of the trombiculid mites, including the specialized setae, without a single reference to this or Grandjean's numerous other publications. It is to be hoped that acarologists generally will realize that it is definitely not in the best interest of acarology for any group of workers to set up a highly formalized and (by virtue of the number who have contributed) ostensibly authoritative system of terminology, either in complete ignorance or in complete disregard of the intensive work of others who have devoted many years of profound study to the morphology of the Acari.

The Trombiculidae are no more than one very small segment of the Acari, and there is no reason why they should be treated differently from other groups, despite the medical importance of a few exceptional members of the family.

In his studies on the Johnstonianidae, the writer has followed the system of nomenclature proposed by Grandjean, first in 1935, and expanded in subsequent years. This will inevitably cause confusion to those familiar with only the terminology proposed by American workers (Wharton *et al.*, 1951). This is not done on the basis of priority, but with the knowledge that a choice must be made between two systems, plus the conviction, based on experience, that the older Grandjean nomenclature is more universally applicable, more fundamental, and, in the final analysis, more logical and comprehensible than the newer highly specialized terminology promulgated by investigators whose experience in the Acari is either primarily or entirely limited to immature stages in a single family.

The following table of equivalents is provided to help in the transposition from one system to the other.

Many examples could be cited to show the inevitable difficulties in the setal terminology which has sprung up during the past few years. One of these difficulties stems from the fact that the terminology has been evolved almost exclusively as an outgrowth of the describing of larvae and shows gross inconsistencies when an attempt is made to apply it to postlarval stages. A second is that related setae are often given quite different names, while some totally unrelated setae are given confusingly similar names.

For instance, the "subterminala" is a specialized seta found near the tip of the tarsus of leg I and the palp. In the adult the ontogenic descendants of the "subterminala" of the palpal tarsus are called "apical setae" or "apical nude setae." The morphological de-

<i>Grandjean System, with Modifications Introduced in This Paper</i>	<i>Terminology Proposed by Various Specialists in the Larval Trombiculidae</i>
solenidion ₁ or s ₁	spur of tarsus I
solenidion ₂ or s ₂	spur of tarsus II
solenidion ₃ or s ₃	microfemorala, microgenuala, microtibiala (in part)
solenidion ₄ or s ₄	microtibiala (in part)
solenidion of palpal tarsus.....	spur of palpal tarsus
dorsal eupathid of larval tarsi, or "eupathid at 0.71d" (etc.)..	subterminala
distinventral eupathid of larval tarsi, or "eupathid at 0.92pv" (etc.).....	pretarsala (a misnomer) (no acceptable equivalent)
eupathidia of various segments of adult legs.....	parasubterminala (no equivalent)
companion seta of dorsal eupathid of larva.....	(no equivalent)
companion seta of s ₁ or s ₂ of larva.....	(no equivalent)
companion seta of s ₄ of larva.....	subterminala of palpal tarsus of larva
eupathidia of palpal tarsus of larva.....	apical setae or apical nude setae (no equivalent)
eupathidia of palpal tarsus of adult.....	microgenuala, microtibiala
supracoxal setae.....	microspur of tarsus I
vestigial setae.....	microspur of tarsus II
famulus ₁	mastifemorala, mastitibiala, mastitarsala
famulus ₂	
bothridia.....	

scendants of the "subterminala" of the tarsi of the legs are simply called "nude setae" in the few cases in which they have been mentioned. "Nude setae" is apparently a catch-all category which includes famuli, solenidia, eupathidia, vestigial setae, and occasional normal setae which lack barbs. What is equally remarkable is that setae which are obviously morphological equivalents of the larval "subterminala" show up on segments as far back as the telofemur in adults. The term "subterminala" is therefore totally inappropriate to designate these setae, while the name "nude setae" is ambiguous. These are all eupathidia, and that term can be applied to these setae whether they are in larva or adult, on palp or leg, on tarsus, tibia, patella or telofemur. The "pretarsala" is another eupathid, but it is not on the pretarsus, and moreover its ontogenic counterparts in the adult are frequently indistinguishable from those of the "subterminala." The eupathidia are a particular type of seta found in a variety of places, and it seems unnecessary to apply special names to them.

The "genuala" and "tibiala" are both solenidia, and nearly all "tibiala" are structurally identical with all "genuala." Why, then,

should these be given different names, just because they are on different segments of the legs? Why not also apply separate names to the normal setae found on the separate segments of the legs? It can also be pointed out that certain of the larval "tibiala" have counterparts which appear on the tarsus of the adult. The "spurs" are also solenidia, although differing structurally from the other solenidia of the legs. Giving the various types of solenidia different names, confusingly parallel in construction in some cases to the names of entirely different types of setae (femorala, genuala, tibiala, microtibiala, microgenuala, microtarsala, subterminala, pretarsala) complicates the picture unnecessarily. What is worse, it obscures the fundamental relationships and true differences between the setal types. These have been very adequately outlined by Grandjean, whose works are to be recommended most highly to any who have not yet read them.

During the course of the present studies it was found necessary to elaborate somewhat upon the solenidial classification in the Parasitengona. The Johnstonianidae, possibly more than any other family, show clearly the multiplicity of form of these setae, but at the

same time they also emphasize the essential unity of the various types. This will be more fully treated below.

The Solenidia of the Legs

The terrestrial Parasitengona are remarkable for the variety of solenidia developed on the legs. These must have considerable physiological, evolutionary and systematic significance, although it will require many years of work before these are understood. At the present time there are few unequivocal statements which can be made regarding the form and distribution of the various types, because of the complexity of study, and the lack of comparative studies in the several families. However, the most significant findings that have emerged from the present study are outlined here to provide at least a tentative beginning toward the eventual comprehensive understanding of the morphology of these important organs. In all cases the statements should be interpreted as applying most immediately to the Johnstonianidae, although exploratory studies in other families show clearly that their applicability extends far beyond the family momentarily under consideration.

1. The several types of solenidia are constant within a given genus. Interspecific variations are found in the numbers of a given type on specific segments of the legs, or differences in the position of certain highly characteristic types, or in minor but constant differences in form of a single type.

2. All species studied have at least three types of solenidia, while most have four. A decrease in number of types comes about through convergence of form rather than through deletion of one or another group of solenidia.

3. The different types of solenidia are recognizable in larval, nymphal and adult stages, although some apparent cases of divergence or convergence in form of two similar types have been observed in the transition from larva to adult.

4. As a general rule the solenidia of tarsi I and II of the larvae are different from each other and from the solenidia of the more proximal segments of the legs. Tarsus III does not possess solenidia in the larvae of the Johnstonianidae. Tarsi I and II in the larva always bear a single dorsal solenidion, usually without, rarely with a companion seta. The tarsal solenidia are thicker and more strongly marked with spiral or annular structure than those of the more proximal segments.

5. For purposes of convenience in differentiating the types and brevity in designating the types, the solenidia of tarsi I and II of the larvae and their ontogenic descendants in nymphs and adults, where these can be distinguished, will be designated solenidia₁ and solenidia₂, or s_1 and s_2 respectively. Where s_1 and s_2 cannot be differentiated this will be interpreted as a convergence in form so that s_1 and s_2 are indistinguishable. The reader should recognize that this is not the only interpretation that is available, but it should be equally apparent that if a different sequence of symbols is used each time a decrease in solenidial types is observed, the description of this change becomes unduly difficult. It is far simpler to utilize a single sequence of symbols which is applicable to the majority of cases and to fit observed exceptions to this rather than to evolve a new system to fit each particular variation. Further justification for this is found in the fact that the larvae of all species of Johnstonianidae studied so far, as well as all species of Trombidiidae and Trombiculidae which have been checked, have four types of solenidia on the legs which can be differentiated by either form or position, or usually both. The apparent reductions in solenidial types which have been observed to date consist entirely of convergences in setal types in the development from larva to adult.

6. The designation solenidion₃ or s_3 will be reserved for the type found on the proximal segments of the legs, with the exception of s_4 (see below). Even in cases in which s_1 and s_2 might be indistinguishable morphologic-

ally, so that they effectively comprise a single type, the designation s_3 will still be applied as indicated here, for the reason explained immediately above. Solenidia₃ are generally small, very slender, usually but not always lacking internal structure, and are probably invariably the most numerous type in both larval and postlarval stages.

7. In the larvae of all genera studied to date, a fourth type of solenidion is found dorsally on tibia I and will be designated solenidion₄ or s_4 . This is a little larger than s_3 and typically shows some degree of internal structure. It may be intermediate in size to s_3 and s_1 , and usually the distinction between these is considerably more difficult to appreciate in the adult than in the larva. The identification of s_4 is also confused by those cases in which a solenidion very similar to s_1 is found on tibia I in the position normally occupied by s_4 . This is true of *Lassenia*, new genus, in which one distidorsal seta of tibia I has a companion seta and is of the same form as s_1 . This might be interpreted either as a case in which s_1 is found on the tibia, or, as the writer has done, a case in which there is a strong convergence between s_1 and s_4 . The situation is in no way simplified by the fact that throughout the protonymphal and deutonymphal instars of *Lassenia* there is a progressive diminution of this seta so that in the adult apparently only one type of solenidion can be found on tibia I with any degree of certainty—namely s_3 .

8. As can be seen from the foregoing, the differences between solenidial types may be very marked, or there may be convergences which make interpretation of types exceedingly complex at times. However, we can only go so far in simplifying a complex situation, and perhaps the more surprising thing would be to find that all patterns of solenidial variation in the Johnstonianidae or the terrestrial Parasitengona in general could be smoothly fitted into a single scheme. If we fail to achieve this, we may yet succeed, if in failing we discover the reason for doing so.

The Johnstonianidae are a very fortunate group for studying the morphology of the solenidia, because in most genera the four types can be recognized not only in the larva but in the adult as well. This is especially so since s_2 in *Centrotrombidium*, *Diplothrombium*, and *Johnstoniana*, are of very unusual form and readily distinguishable from s_1 and s_4 . In *Lassenia* larvae, s_1 and s_2 are of similar form and size, differing chiefly in the presence of a companion seta at the base of s_1 . However, the companion seta is never retained in the postlarval instars, so this difference cannot be utilized in differentiating the two types of solenidia in the nymphs and adults. Consequently in this genus it is all but impossible to differentiate between s_1 and s_2 in the adult; moreover this situation extends to s_4 as well, as pointed out above.

One interesting feature of s_2 is that this type, while it appears first on tarsus II of the larva, is actually more abundant on tarsus I in the adult than it is on tarsus II. In other words they arise *de novo* on tarsus I in the postlarval stages, and in greater numbers there than in the site at which s_2 originally appeared in the larva. This is true in all genera (except possibly *Lassenia* in which the true state of affairs has not been ascertained because of the convergence in solenidial types in the postlarval stages).

The Eupathidia of the Legs of the Larva

Each of the four genera studied by the writer has a characteristic arrangement of the larval eupathidia. In all genera, the eupathidia are confined to the tarsi, and in all cases tarsus I has two eupathidia. Variations are found in the number of eupathidia on II and III, and in the presence or absence of companion setae. Both *Centrotrombidium* and *Johnstoniana* have eupathidial formulae of (2-1-0), but only in *Johnstoniana* is there a companion seta at the base of the dorsal eupathid. *Diplothrombium* has a formula of (2-1-0) in both species seen by the writer, but there is no companion seta with any of the eupathidia. In the two

species of *Lassenia* known in the larval stage, the eupathidial formula is (2-2-1), and the dorsal eupathid of both tarsi I and II in both species has a basal companion seta. Only in *Lassenia* is there a eupathid on tarsus III, and this is subterminal in position.

The Eupathidia of the Legs of the Adult

Here we find differences in the distribution of the eupathidia on the various segments of the legs, with *Centrotrombidium* having the eupathidia confined to the tarsi, and tarsus IV typically with only one eupathid. In the other three genera, eupathidia are found on all segments beyond the basifemur. In all genera but *Lassenia* there is a subterminal eupathid on tarsus IV. In *Centrotrombidium* this lies at about $0.85a$ to av , in *Diplothrombium* at 0.90 to $0.95v$ and in *Johnstoniana* at $0.91pv$. Specific differences are found in the distribution of the eupathidia on the individual segments of the legs. One interesting example is found in the genus *Lassenia* in which ventral eupathidia are found on the telofemur, patella and tibia of leg I, whereas in *L. lasseni* all of the eupathidia are dorsal or marginal in position.

The Companion Setae

Companion setae are found only in the larvae of some genera, and are seemingly universally absent in the postlarval stages of all of the Parasitengona. Neither *Centrotrombidium* nor *Diplothrombium* has any companion setae in either larval or postlarval stages. *Johnstoniana* larvae have one companion seta associated with the dorsal eupathid of tarsus I, while in *Lassenia* there are four companion setae. These are associated with s_4 of tibia I, s_1 of tarsus I, and the dorsal eupathid of both tarsi I and II.

The Famulus

The famulus is the most constant of all the specialized setae in the Johnstonianidae. A famulus is present on both tarsi I and II in all genera, in both larval and adult stages.

Variations in the position of the famulus often provide very useful specific characters (as in *Centrotrombidium*, for example), although a noticeable degree of variation in the position of the famulus is observed in some species. In others the famulus appears to show extremely little variation in position.

The Supracoxal Setae

The situation here is not certain, although it is probable that these are not as variable in their number and distribution as are the setae on the ventral surface of the coxa. Supracoxal setae, when present, are found both on the dorsal surface of the coxal portion of the gnathosoma, and on the dorsal surface of the coxae of leg I. This appears to be a general rule in the Parasitengona. In the Johnstonianidae, supracoxal setae are absent in all but the genus *Lassenia*. In the known species of this genus supracoxal setae are present on the dorsal surface of the gnathosoma as well as on the dorsal surface of coxa I. The distribution of supracoxal setae in the adult is always the same as in the larva so far as is known.

The Vestigial Setae

These are the small spikelike setae at the distidorsal end of the patella or tibia of legs I and II (Figs. 91, 194, 215). The distribution of these very often follows that of the supracoxal setae, that is, when the latter are present vestigial setae are also present. This is true in both *Centrotrombidium* and *Diplothrombium* in which none of the described species has either supracoxal or vestigial setae. In *Lassenia lasseni* and *L. spinifera*, vestigial setae are found on patella I and II and on tibia I; likewise supracoxal setae are present on the palpi and leg I. In *Johnstoniana latiscuta* an intermediate condition is found; for although there are no supracoxal setae, vestigial setae are present on patella I and II but are absent from all tibiae. As in the case of the supracoxal setae the distribution of the vestigial setae is identical in larva and adult in all species studied. The distribution of the supracoxal and vestigial

DISTRIBUTION OF COXAL, SUPRACOXAL AND VESTIGIAL SETAE (LARVAE)

	No. Coxal Setae	Supracoxal Setae		Vestigial Setae			
		Palpi	I	I		II	
				pa	ti	pa	ti
<i>J. latiscuta</i>	2-1-1	0	0	1	0	1	0
<i>D. monoense</i>	2-1-1	0	0	0	0	0	0
<i>D. cascadiense</i>	2-1-1	0	0	0	0	0	0
<i>C. distans</i>	2-1-1	0	0	0	0	0	0
<i>L. lassenii</i>	2-2-3	1	1	1	1	1	0
<i>L. scutellata</i>	2-1-2	1	1	1	1	1	0

setae in the larvae of the known Johnstonianidae is summarized in the table given above. Although this table is based on the larvae, the figures would be identical for the adults except for the number of coxal setae.

The Specialized Setae of the Palpal Tarsus of the Larva

One constant feature in the chaetotaxy of the palpal tarsus of the larva is the presence of a single solenidion on the posterior aspect of the basal one-third of the segment. The tarsal eupathidia exhibit one interesting variant. In both species of *Lassenia* these are of typical eupathidiform structure, tubular throughout, and relatively thin-walled. Moreover, the terminal eupathid is always at the very end of the segment (Fig. 253). In the other three genera, the end of the tarsus always extends beyond the insertion of the most distal seta. Moreover, none of the tarsal setae in larvae of these three genera are typically eupathidiform, although it may be presumed that some of the terminal ones are homologous with the typical eupathidia of *Lassenia*. These subterminal setae are strongly hemipectinate, and in *Centrotrombidium distans*, the one nearest the end of the segment is also flattened and expanded (Figs. 28, 29, 92, 176).

The Specialized Setae of the Palpal Tarsus of the Adult

While in the larva the solenidion on the posterior surface of the palpal tarsus is always in the basal one-half or one-third of the segment, in the adult there is a strong tendency

for the solenidion to be displaced toward the distal end. In *Centrotrombidium* it is found at roughly 0.3*p*, in *Diplothrombium* at approximately 0.5*p*, and in *Johnstoniana* and *Lassenia* at approximately 0.8 to 0.9*p*. Paralleling the displacement of the solenidion is an increase in number of eupathidia. Thus, in *Centrotrombidium* there are only 2 terminal eupathidia, but in *Diplothrombium* there are 3 to 5, in *Johnstoniana* about 6, and in *Lassenia* from 6 to 15 depending upon the species and the individual. These trends are not entirely associated with size for although *Diplothrombium micidium* is smaller than *D. monoense*, the number of eupathidia on the tarsus of the palp is greater. Also the tarsus of the palp of *Johnstoniana latiscuta* is larger than that of *Lassenia spinifera*, yet there are only about half as many eupathidia.

DESCRIPTIONS AND KEYS

JOHNSTONIANIDAE new family

DIAGNOSIS: Adults with genital and paragenital sclerites about equally developed, and with three pairs of genital acetabula. Scutum with two pairs of typical sensilla, or with anterior pair greatly modified to absent. Scutum frequently with an anterior spine. Hysterosomal setae typically smooth, borne on individual sclerites. Chelicerae as in Trombididae, with base, movable tarsus, and a delicate dorsal membrane. Palpal tibia without ctenidium, with only the one terminal and one subterminal spiniform setae. Anterior wall of palpal trochanter often fenestrated.

Larvae also with one to two pairs of sensilla on the scutum. Dorsum with setae borne

individually on relatively large sclerites. Intercoxal area with a single pair of setae between III, or devoid of setae; urpore present. Either three or two pairs of rostral setae. Palpal tarsus simple, cylindrical or fusiform. Larvae self-detaching parasites of insects, living in damp to very wet places, and typically with relatively restricted temperature tolerances.

REMARKS: Not all of the variants of the morphological characters of the Parasitengona can be readily utilized in a key, but the more convenient ones are included here. It is interesting that the number of characters available in the larva considerably exceeds the number available in the adult. Actually the discrepancy is not as great as would appear at first glance, since some adult characters are omitted simply because of difficulties in interpreting them.

It appears possible that the Johnstonianidae as defined here will require further reorganization as more forms become known from other parts of the world. At present it is quite evident that there are two distinct groups of genera, the first including *Centrotrombidium*, *Diplothrombium*, *Johnstoniana*, and provisionally *Hirstithrombium* (Johnstonianinae Thor 1935). The second includes *Lassenia* new genus, *Polydiscia* Methlagl 1928, and provisionally *Crossothrombium* Womersley 1939 (*LASSENIINAE* new subfamily). The principal structural differences between the genera are summarized in the formula keys given below.

FORMULA KEY TO GENERA OF

JOHNSTONIANIDAE

Based on Adults

- 1a. Anterior sensilla present, similar in form to the posterior sensilla, and at least one-half as long as the posterior sensilla (Figs. 70, 160).
- 1b. Anterior sensilla present, but considerably modified, less than one-half as long as the posterior sensilla (Figs. 183, 232).
- 1c. Anterior sensilla absent; scutum greatly

reduced, bearing a single pair of sensilla, anterior to which is a single pair of normal setae (Fig. 4).

- 2a. Pregenital tubercle absent.
- 2b. Pregenital tubercle present (Figs. 185, 233).
- 3a. Anal sclerites present (Figs. 18, 105).
- 3b. Anus surrounded only by membranous cuticle containing setae, but no sclerites (Fig. 68).
- 4a. Palpal trochanter with anterior surface containing a distinct oval fenestra (Fig. 61).
- 4b. Distal margin of anterior wall of trochanter deeply incised, but the fenestra is not completely circumscribed (Fig. 201).
- 4c. Anterior wall of trochanter with no trace of a fenestra (Fig. 69).
- 5a. Eupathidia present on leg segments other than the tarsi.
- 5b. Eupathidia confined to the tarsi.
- 6a. Supracoxal setae absent from gnathosoma and coxae I.
- 6b. Supracoxal setae present on gnathosoma and coxa I.
- 7a. Vestigial setae absent from patella and tibia of all legs.
- 7b. Vestigial setae present only on patella I and II, absent from tibiae.
- 7c. Vestigial setae present on patella I and II, and tibia I; absent from tibia II.
- 8a. Solenidia₂ of tarsus I clavate, markedly different in form from s₁.
- 8b. Solenidia₁ and s₂ of tarsus I differing primarily in size; no sharp differences in form.

FORMULA KEY TO GENERA OF

JOHNSTONIANIDAE

Based on Larvae

- 1a. Scutum with four pairs of setae (two pairs of normal setae plus two pairs of sensilla).
- 1b. Scutum with two pairs of setae (one pair of normal setae plus one pair of sensilla).

DISTRIBUTION OF VARIANTS

1	2	3	4	5	6	7	8	JOHNSTONIANINAE
a	a	b	c	a	a	b	a	Johnstoniana
a	a	a	a	a	a	a	a	Diplothrombium
c	a	a	a	b	a	a	a	Centrotrombidium
?	?	?	?	?	?	?	?	Hirstithrombium
								LASSENIINAE
b	b	a	b	a	b	c	b	Lassenia
?	?	?	?	?	?	?	?	Polydiscia
?	?	?	?	?	?	?	?	Crossothrombium

- 2a. Anal sclerites absent.
- 2b. Anal sclerites present, usually with one or more pairs of setae.
- 3a. Protorostral, deutorostral and tritorostral setae all present and setiform or spiniform (deutorostrals often concealed under velum).
- 3b. Deutorostral setae absent or reduced to extremely minute vestiges, tip of rostrum with only two pairs of setae.
- 4a. Palpal trochanter with anterior wall fenestrated.
- 4b. Anterior wall of palpal trochanter not fenestrated.
- 5a. Terminal seta of palpal tarsus eupathidiform (hollow, thin-walled, etc.).
- 5b. Terminal seta of palpal tarsus not typically eupathidiform, but flattened, strongly serrate, or otherwise modified.
- 6a. Tarsal eupathidia 2-1-0.
- 6b. Tarsal eupathidia 2-1-1.
- 6c. Tarsal eupathidia 2-2-1.
- 7a. Dorsal eupathid of tarsus I with companion seta.
- 7b. Dorsal eupathid of tarsus I without companion seta.
- 8a. Dorsal eupathid of tarsus II present, with companion seta.
- 8b. Dorsal eupathid of tarsus II absent.
- 9a. Solenidion₄ of tibia I with companion seta.
- 9b. Solenidion₄ of tibia I without companion seta.
- 10a. Coxal setae with formula 2-1-1.
 - b. Coxal setae with formula 2-1-2 or 2-2-3.

- 11a. Supracoxal setae absent on gnathosoma and coxae I.
- b. Supracoxal setae present on gnathosoma and coxae I.
- 12a. Vestigial setae absent from patella and tibia of all legs.
- b. Vestigial setae present on patella I and II only, absent from tibiae.
- c. Vestigial setae present on patella I and II and tibia I; absent from tibia I.
- 13a. Femora I-III of all legs completely divided into basifemur and telofemur (six free segments beyond the coxae).
- b. Femora I-III undivided (only five segments beyond the coxae).
- c. Femora I-III partially divided, the synarthrodial membrane well formed ventrally, but absent dorsally (Figs. 87, 91, 93).
- 14a. Tarsi with two unequal claws on all legs, the anterior claw more erect than the posterior.
- b. Tarsi with three claws on all legs, the anterior and posterior claws equal, the median (axial) claw more slender, erect.
- 15a. Intercoxal area with a pair of setae between III.
- b. Intercoxal area without a pair of setae between III.

JOHNSTONIANINAE Thor 1935

DIAGNOSIS: Adults with either one or two pairs of sensilla on scutum. Pregenital tubercle absent. Solenidia₂ typically clavate, differing appreciably from s₁. Supracoxal setae absent from gnathosoma and coxa I in both larva and adult. Larvae without anal sclerites; usually with a single pair of setae in the inter-

DISTRIBUTION OF VARIANTS															
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	JOHNSTONIANINAE
a	a	b	b	b	a	a	b	b	a	a	b	c	a	?	Johnstoniana
a	a	b	a	b	a	b	b	b	a	a	a	a	a	a	Diplothrombium
b	a	b	a	b	a	b	b	b	a	a	a	a	a	a	Centrotrombidium
?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	Hirstithrombium
															LASSENIINAE
a	b	a	b	a	c	a	a	a	b	b	c	b	b	b	Lassenia
a	b	?	?	a	?	?	?	?	b	?	?	a	b	b	Polydiscia
?	?	?	?	?	?	?	?	?	?	?	?	?	?	?	Crossothrombium

coxal area between coxae III. Deutorostr al setae absent; terminal seta of palpal tarsus not eupathidiform. Tarsi each with two claws.

Centrotrombidium Kramer 1896

ADULT: Length of idiosoma 500 to 1200 μ in known species. Scutum of both adult and larva bearing a single pair of clavate sensilla and anterior to these, a single pair of elongate normal setae; anterior sensilla absent. Anterior end of scutum drawn out into a long spine. Ocular plates bearing two refractile corneae each, or one refractile cornea and a non-refractile lobe. Dorsal hysterosomal setae borne on individual sclerites, no large plates present. Anal sclerites well developed. Supracoxal setae of legs and gnathosoma and vestigial setae of legs completely absent in both larva and adult. Trochanter of palp with anterior wall fenestrated in both larva and adult. Palpal tibia with a single subterminal spiniform seta; tarsus with one solenidion and usually only two eupathidia. Solenidia of legs of four recognizably different types; tarsi I and II with one or more clavate solenidia₃ on posterior aspect. Famulus of tarsi I and II located in distal one-fifth of segment, on posterior membrane of claw fossa. Eupathidia of legs confined to tarsi.

LARVA: In addition to those which are common to both larva and adult, the larva has the following characteristics. Ocular plates distinctly bicornate. Anal sclerites absent. Velum small, simple; deutorostr al setae absent, or represented by vestigial structures. Base of gnathosoma without posterorostr al setae. Femur of all legs divided. A single

clavate solenidion₂ on tarsus II, s₁ of tarsus I curved, rodlike. Tarsus of all legs with two simple, smooth claws.

REMARKS: At present only five or six species are known, including the three new ones described here. *C. schneideri* Kramer 1896 is rather widely distributed in Europe, and *C. australasiae* is known from Australia. *C. misellum* (Berlese), from Mexico, is provisionally placed in this genus. The genus is probably cosmopolitan. The adults are small, brown mites, rather inconspicuous, found crawling over wet ground along the margins of streams and ponds. The writer has also collected them in estuarine habitats. The larvae are probably parasitic upon various small arthropods but nothing is known about host relationships. The known species can be separated on the basis of the diagnostic formulae given below.

FORMULA KEY TO ADULTS OF CENTROTROMBIDIUM

- 1a. Sensilla of scutum with distal portion nearly spherical, length of stalk less than twice the diameter of the sphere, and less than the distance between the sensillar alveoli.
- 1b. Distal portion of sensillum clavate or pyriform, stalk very long, at least twice as long as the expanded tip, and considerably longer than the distance between the alveoli.
- 1c. Sensilla not swollen, but slender through out.
- 2a. Dorsal hysterosomal setae relatively long-slender, tapering uniformly to a very fine point (Figs. 19, 22).

- 2b. Dorsal hysterosomal setae appreciably stouter and of more nearly uniform diameter throughout, reaching only a little beyond the margin of the setigerous sclerites (Fig. 64).
- 3a. Ocular plates with only one distinct cornea, plus a non-refractile posterior lobe (Fig. 53).
- 3b. Ocular plates with two distinct corneae.
- 4a. Tarsus I with only two (exceptionally 1 or 3) clavate solenidia₃ on posterior membrane of claw fossa (Fig. 48).
- 4b. Tarsus I typically with more than two clavate solenidia here.
- 5a. Famulus of tarsus II very close to alveolus of clavate solenidion₃, removed from it by a distance no greater than the length of the solenidion, and with no other setal alveoli in the intervening distance (Fig. 49).
- 5b. Famulus of tarsus II removed from alveolus of clavate solenidion₃ by a distance greater than the length of the solenidion; one or two normal setae in intervening distance (Figs. 6, 8).
- 6. Range in idiosomal length, by sex, where known.

Since the larva of only *C. distans* is known, no key to larvae can be given. The above key does not include all the characters of probable specific value in the genus, but they are the more important ones. Others include the difference in form of the posterior end of the scutum, which is important in differentiating *C. distans* and *C. approximatum*, but this is difficult to state in words. Womersley (1942, p. 171, fig. 2c) showed no anterior spine on the scutum of *C. australasiae* Womersley 1942. It is likely that this was drawn from an un-

dissected specimen, and in such specimens the scutal spine often projects straight downward so that it cannot be seen in dorsal view. Whether or not this is the case in Womersley's species is not known, but the possibility should certainly be kept in mind upon comparing this with other species of the genus.


Centrotrombidium distans new species

FEMALE: Idiosoma (Fig. 1) 897–988 μ long, 702–728 μ wide (two ovigerous specimens). Scutum (Fig. 4) drawn out into a long anterior spine, and with a single pair of capitate sensilla, anterior to which is a pair of stout, long, smooth to faintly roughened setae; anterior sensilla absent. Crista metopica extending for most of length of scutum. Ocular plates very small, bearing only one prominent cornea (the anterior one) and an angular process (Fig. 25). Cuticle of ocular plate outside of cornea minutely tuberculate. Propodosoma between ocular plate and scutum with about seven setae, all of which lie at or anterior to the level of the sensilla. Dorsal and lateral body setae (Fig. 19) arising from alveoli borne on prominent raised sclerites, shaft sharply deflexed near base of seta. Membranous cuticle between sclerites with no visible markings except for a few faint parallel striae seen in some parts of the body. Subcuticular reticular layer a network of fine fibrils.

Coxae I and II (Fig. 3) with only about 20 to 25 smooth setae each; pars medialis coxae with about 5 setae; supracoxal seta absent. One small apodeme behind coxa II, very feebly developed. Coxae III and IV also with about 20 to 25 setae (Fig. 5). Ventral body setae like those of dorsal and lateral surfaces. Genital sclerites (Fig. 10) with 9 to 12 smooth

DISTRIBUTION OF VARIANTS

1	2	3	4	5	6	
a	a	a	a	b	♀ 897– 988 μ	<i>distans</i> n. sp.
a	a	a	a	a	♀ 988–1092 μ	<i>approximatum</i> n. sp.
a	a	ab	a	b	♀ 962 μ	<i>hadroseta</i> n. sp.
a	a	b	?	?	525 μ	<i>schneideri</i> Kramer 1896
c	b?	?	?	?	800 μ	<i>misellum</i> (Berl.) 1918
b	?	b	b?	?	1020 μ	<i>australasiae</i> Wom. 1942



FIGS. 1 - 9. *Centrotrombidium distans* n. sp.: 1, dorsum, female; 2, tip of rostrum showing tritrostral seta and presumed deutrostral rudiment; 3, coxae I and II, female; 4, propodosoma, female; 5, coxae III and IV, female; 6, tarsus I, female, posterior; 7, cheliceral tarsus, female; 8, tarsus II, male, posterodorsal; 9, tarsus I, posterior, female.

setae in a single row, paragenital sclerites with 15 to 18 smooth setae. Three pairs of genital acetabula. Anus (Fig. 18) flanked by two crescentic sclerites bearing 6 to 8 smooth setae each. Base of gnathosoma and rostrum as described for *C. approximatum*, new species. Chelicerae also as in the following species, except that the teeth on the chelicerae appear to be significantly heavier (compare Figs. 7, 52). In the holotype female there was a seta at the base of the dorsal membrane of one chelicera, and an indication of at least the alveolus and a short shaft on the other chelicera (Fig. 24). However, no such seta could be found on the male collected at the same locality, nor on either the male or female of *C. approximatum*. This is an interesting anomaly, suggestive of the condition normally found in the Parasitiformes, and many of the Eleutherengona in which a seta is frequently found in this position in normal individuals. Number, form, and arrangement of setae on all segments of the palp identical with that of the holotype of *C. approximatum* (Figs. 58, 59, 61).

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae, m = many).


Vestigial setae absent on all segments of all legs, and eupathidia present only on the tarsi. Tarsus I greatly swollen, height/length 0.50 to 0.56 (two specimens), claw fossa beginning at 0.64 to 0.65 and extending to end of tarsus. Typically with two clavate solenidia₂ on the posterior aspect of tarsus I, the first of these at about 0.65 to 0.68*pd*, the second at 0.70 to 0.79*pd* (Fig. 9). In some cases *s*₂ have a faintly annulate structure; in others they appear smooth. The peculiar seta with the

fusiform distal half seen disti-ventrally to the solenidia₂ in Figure 9 is an anomaly. Famulus spikelike, located at 0.90 to 0.93*pd*, arising from a distinctly vesicular alveolus. Tarsus I also with numerous *s*₁ and eupathidia, the latter extending around the ventral and distal margin of the tarsus from 0.43*v* to 0.80*d*. The eupathidia have a characteristically geniculate terminal filament visible under oil (not shown in figure). Some tarsi I have only one or as many as three solenidia₂, but two is the normal number. Tarsus II with a single clavate solenidia₂ near the basal end of the claw fossa (Figs. 6, 8) at about 0.64*pd*, and a peglike famulus at 0.85*pd*, borne on a distinctly vesicular alveolus. Tarsi III and IV with two and one eupathidia each in the material studied. *S*₂ highly distinctive (Figs. 15, 16), but *s*₄ (Fig. 13) occasionally intergrading with *s*₁ and *s*₃ so that it is difficult to assign certain setae to one or another type.

MALE: Body 754 to 858 μ long (average 806 μ , three specimens). In general resembling female except for structure of genital area (Fig. 21). Genital sclerites with 6 to 9 setae in a single row, paragenital sclerites with 12 to 18 setae in a single row. Three pairs of genital acetabula. An interior circlet of about 12 setae can be seen in some favorable specimens, but are not shown in the figure. Penis very small, inconspicuous. Anal sclerites as in female.

LARVA: Idiosoma (Fig. 27) 152–167 μ long. Scutum somewhat triangular, greatly reduced (Fig. 32), bearing only the posterior pair of sensilla, anterior to which is a pair of stiff, hemipectinate setae. Between the latter a very short, rudimentary crista metopica. Sensilla capitate, the spherical distal portion crumpled in some mounted specimens (artifact). Ocular

	tr	bf	tf	pa		ti		ta				e	f
	n	n	<i>s</i> ₃	<i>s</i> ₃	<i>s</i> ₄	<i>s</i> ₃	<i>s</i> ₄	<i>s</i> ₁	<i>s</i> ₂	<i>s</i> ₃	<i>s</i> ₄	m	1
I	4–5	15	6–10	31	2–3	28–35	3	15	2	0	4?		
II	4	11	3–7	16	0	17	2	6–8	1	0	0	2	1
III	5	11	3–4	8–12	0	10–14	0	0	0	0	2	2	0
IV	8	9	4	9	0	8	0	0	0	0	1	1	0



FIGS. 10-30. *Centrotrombidium distans* n. sp.: 10, genital opening, female; 11, solenidion, patella I; 12, s₃, femur II; 13, s₄, tarsus II; 14, s₁, tarsus I; 15, s₂, tarsus I; 16, s₂, tarsus II (all from female); 17, gnathosoma, female; 18, anus, female; 19, dorsal hysterosomal setae, side view; 20, same, top view; 21, genital area, male; 22, marginal body seta, female; 23, rostrum, female, lateral; 24, chelicera showing abnormal seta, female; 25, ocular plates, female; 26, venter, larva; 27, dorsum, larva; 28, palpal tibia and tarsus, larva, posterior; 29, terminal seta of palpal tarsus, larva; 30, tibia and tarsus of palp, larva, anterior.

plates bicornate, simple in form. Dorsal and marginal setae numbering 24 in four rows of 6 each, each seta hemipectinate and borne on a separate sclerite.

Coxa I with one pectinate seta laterally and another in the medial angle (Fig. 26); supra-coxal seta absent. Medial portion of coxa I very indistinct, differentiated only by the absence of striae. Coxa III with a single bifurcate seta. Intercoxal area with a single pair of smooth setae, inserted on a pair of smooth, slender sclerites between III. Fifteen post-coxal setae on each side, borne on individual sclerites surrounding the anus.

Base of gnathosoma (Fig. 45) with one pair of simple smooth setae (tritorostrals?), deutorostral setae vestigial, visible only under oil immersion; protorostrals small, simple. Velum oval in outline, reticular in appearance. Chelicerae small, compact (Fig. 44), digitus fixus very small, pointed, tarsus not in favorable position for study, but apparently with one dorsal tooth proximal to the tip. Palpi (Fig. 33) five-segmented, trochanter lacking setae, femur and patella each with a single dorsal seta with a few barbs. Palpal tibia with three simple setae and the unidentate spurlike seta which is inserted in the end of a truncate projection of the tibia. Palpal tarsus (Figs. 28, 30) with solenidion at 0.10*p*, with only the faintest indication of annuli. A curved, hemipectinate seta at 0.50*av* which extends along ventral margin of tarsus, somewhat obscuring the very fine tip. A very heavy seta at 0.71*d*, terminal portion flattened, spatulate, the margin pectinate. Tip of tarsus forming a sharp pointed spine extending well beyond insertion of spatulate seta. Otherwise with five setae which are smooth, or which bear from one to four pectinations; typical eupathidia absent.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae).

Tibia I with one *s*₃ and one *s*₄ dorsally, II and III each with one *s*₃, and with a ventral keel which is best developed on III. The solenidia on the first five segments of the legs have no perceptible spiral structure.

Tarsus I (Fig. 31) with *s*₁ at 0.47*pd*, a smooth eupathid at 0.75*pd*, and a heavy, smooth eupathid at 0.87*v*. A smooth normal seta at 0.80*d*. Otherwise with 25 hemipectinate normal setae. Claws as described for II. Tarsus II (Figs. 34, 35) with a spikelike famulus at 0.68*pd*, and a clavate solenidion, at 0.67*d*, the stalk sometimes showing distinct spiral structure. A long, smooth, normal seta at 0.81*d* and a eupathid at 0.80*pv*, otherwise with 21 hemipectinate normal setae. Claws two in number, the anterior one smoothly curved, scythe shaped, the posterior one with similar taper and thickness, but with a sharp 90 degree flexure between basal and middle thirds. Tarsus III (Fig. 46) with a smooth normal seta at 0.65*d*; otherwise with 14 normal hemipectinate setae. Tarsal claw as on I and II.

TYPE LOCALITY: Tule Lake, Siskiyou County, California (holotype female). Common along alkali-encrusted shore of lake, on mud overgrown with grasses and other plants. Types in author's collection.

REMARKS: This species is widely distributed throughout the western United States, living in marshy situations. It has also been collected at a point about 4 miles north of Tonasket, Okanogan County, Washington, on the borders of small ponds on the Okanogan River. These ponds are permanent, and are usually flooded each year. A second species, *C. ap-*

	tr	bf	tf		pa		ti			ta				
	n	n	s ₃	n	s ₃	n	s ₃	s ₄	n	s ₁	s ₂	e	f	n
I	1	1	1	5	4	5	1	1	4	1	0	2	1	26
II	1	2	1	4	2	4	1	0	4	0	1	1	1	22
III	1	2	1	4	2	4	1	0	4	0	0	0	0	15

proximatum n. sp. was found at the latter locality. This is very similar to *C. distans* but differs in at least three consistent respects. Because of the high degree of variation found in the Johnstonianidae and the terrestrial Parasitengona in general, it is often difficult to assess the significance of small variations between individuals. However, all individuals of *Centrotrombidium* seen by the writer fall into one of two groups depending upon the degree of separation between the clavate solenidion₂ and the famulus on tarsus II (Figs. 6, 49). In both males and females of *C. distans* these are widely separated as described above; whereas in the others the clavate solenidion and famulus are very close together. Moreover, the famulus in *C. distans* is borne on a protruding vesicular base while that of the other form arises from an alveolus which is set at or below the general surface of the cuticle. The second arrangement has been found in both males and females so that sexual differences are not involved. A second difference is in the number of solenidia on the dorsal surface of tibia I. These segments are approximately the same size in the two forms, but in *C. distans* there are approximately 34 to 41 solenidia; whereas in the second form there are only about 20. A third difference, in the form of the scutum, is discussed in the description of *C. approximatum*.


Centrotrombidium approximatum new species

FEMALE: Very similar to *Centrotrombidium distans* new species in size and general appearance. Length 988 to 1092 μ to tip of scutal spine, width 728 to 884 μ , length/width 1.23 to 1.35; average 1050 μ long, 801 μ wide, length/width 1.31 (five ovigerous females). Scutum slightly larger than in *C. distans*, with the portion behind the area sensilligera of nearly uniform width throughout in contrast with the expanded form of this part in *C. distans*. Crista metopica well developed, the portion between the area sensilligera and the scutal spine being somewhat longer generally


than in *C. distans*, but this is a variable feature. Posterior sensilla spherical distally, anterior sensilla absent; scutum with only one pair of setae in addition to the sensilla. With 2 to 8 setae between the scutum and each ocular plate (average 4.5 setae, six specimens). Ocular plates appreciably larger than in *C. distans* (Fig. 53) bearing a single cornea laterally and a large lobe medially. The medial lobe is punctate like the rest of the ocular plate, and apparently is never tuberculate as it frequently is in *C. distans*. Dorsal and marginal body setae apparently slightly longer than in *C. distans*, each borne on a separate sclerite, but these sclerites are more nearly hemispherical, and noticeably less elevated than in the preceding species (Fig. 57). Membranous cuticle of dorsum smooth, subcutaneous reticular layer as in *C. distans*.

Coxae I and II with 27 and 26 smooth, slender setae each, pars medialis coxae with 5 setae (one specimen). Coxal ring I incomplete dorsally, supracoxal setae absent; coxal ring II complete. Both I and II with a large articulating process in anterior portion of coxal ring (Fig. 54). Intercoxal area of I and II with 9 to 20 smooth, simple setae, each borne on an individual sclerite; no apodemes behind II. Coxae III and IV with 27 and 25 smooth slender setae respectively in specimen counted, coxal rings complete dorsally; also with a prominent articular process on the anterior half of the coxal ring. Genital and paragenital sclerites as in *C. distans* (Fig. 10), genital sclerites with a single row of 9 to 11 setae, paragenital sclerites with a single to double row of 12 to 17 setae. Three pairs of genital acetabula. Anal sclerites also as in *C. distans*, with 6 to 10 setae on each side.

Base of gnathosoma essentially as in *C. distans* but with a larger number of setae (6 to 10). Velum directed anteriorly, protorostral setae of same form as in *C. distans*; a pair of ventrally directed setae distiventrally on rostrum as in the case of *C. distans*. It appears probable that these are the tritrostral setae, for between these and the protorostrals is a



FIGS. 31-46. *Centrotrombidium distans* n. sp., larva: 31, tarsus I; 32, scutum (sensillum crumpled); 33, palp, posterior; 34, famulus and solenidion₂, tarsus II; 35, tarsus II; 36, solenidion₄, tibia I; 37, s_3 , patella III; 38, s_3 , femur III; 39, s_1 , tarsus I; 40, s_2 , tarsus II; 41, leg I; 42, leg II; 43, leg III; 44, chelicerae, dorsal; 45, gnathosoma, ventral; 46, tarsus III.



FIGS. 47-57. *Centrotrombidium approximatum* n. sp.: 47, propodosoma, male; 48, tarsus I, female, posterior; 49, tarsus II, male, posterior; 50, legs I and II; 51, legs III and IV, female; 52, cheliceral tarsus, female; 53, ocular plates, female to left, male to right; 54, coxae I and II, female; 55, coxae III and IV, female; 56, hysterosomal setae, female, from above; 57, hysterosomal setae, side view.

pair of minute structures like those found in *C. distans* which may represent the vestiges of the deutorostral setae. Including the proto-rostrals and tritrostrals the rostrum bears 10 to 16 setae. Chelicerae (Fig. 60) of standard form, tarsus heavily sclerotized, appearing smooth at low magnification but with 20 to 30 extremely minute teeth visible at magnifications of 250 x or higher. Trochanter of palp (Fig. 61) with fenestral membrane on anterior aspect, setae absent; femur with about 12 smooth straight slender setae. Two of the normal setae on the femur of the palp appear to occupy a characteristic posterior position, isolated from the remaining 10 setae of the segment. Patella with about 9 setae, tibia with 8 smooth slender setae plus the large clawlike unidentate distal seta, and a shorter spiniform seta near its base. Tarsus of palp with three rather heavy normal setae along dorsal margin (Figs. 58, 59), two heavy eupathids distally, and a single solenidion posteriorly.

MALE: Resembling female in all essential respects noted above except for structure of genital area. Genital sclerites and paragenital sclerites as in female, but with slightly fewer setae in most cases. Penis of rather characteristic form, with three elongate apodemes. Genital acetabula numbering three pairs. Anal sclerites also as in female.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae, m = many).

Basifemur I with about 20 to 25 normal setae. Vestigial setae absent; eupathidia found only on the tarsi. Tarsus I with famulus at 0.89pd and with two solenidia₂ on posterior aspect (only one in some cases). Tarsus II with s₂ at about 0.64pd, and with a spikelike famulus at 0.64 to 0.73pd. Eupathidia extend-

ing from 0.55v to 0.95d. Eupathidia of tarsus I extending from 0.49v to 0.85d. Tarsus I also with numerous s₁ and a number of smaller solenidia intermediate in size and form between s₁ and s₃, and assigned in the table to type s₄. There appears to be fairly complete intergradation between s₄ and s₁ as in the case of *C. distans*.


TYPE LOCALITY: About 4 miles north of Tonasket, Okanogan County, Washington, from border of small permanent pond. August 14, 1952. Collected by the writer. This species has not been found elsewhere, and may be a more northerly form, whose principal range extends into British Columbia.

REMARKS: This species is very similar to *Centrotrombidium distans* new species, but differs consistently in the close approximation of s₂ and the famulus of tarsus II (hence the specific name). It also differs in having considerably fewer solenidia on the legs (approximately 125 compared with 200 in *C. distans*). The difference is largely due to the discrepancy in the number of s₃ (about 80 in *C. approximatum*, compared with 160 in *C. distans*); the other solenidial types are present in approximately equal numbers in the two species. Other differences are in the form of the posterior end of the scutum, and the form of the ocular plate as indicated in the foregoing description.

Centrotrombidium hadroseta new species

FEMALE: A single specimen of this form was found at the type locality of *C. approximatum*. It is closely related to *C. distans* from which it differs in only very minor respects. Body of same form as *C. distans*, 962 μ long, 650 μ wide, length/width 1.48. Scutum (Fig. 63) essentially as in *C. distans*, except that the setae in front of the sensilla appear to be

	tr	bf	tf	pa		ti		ta					
	n	n	s ₃	s ₃	s ₄	s ₃	s ₄	s ₁	s ₂	s ₃	s ₄	e	f
I	4-5	23	11	31	2-3	7-9	1	11	2,1	0	7	m	1
II	4-5	15-16	6	10	1	6	1	11	1	0	0	m	1
III	6	11-13	4	8	2	4	1	0	0	2	3	8	0
IV	12	9-10	4	6	0	4-5	0	0	0	1	1,2	1,0	0



FIGS. 58-61. *Centrotrombidium approximatum* n. sp.: 58, tibia and tarsus of palp, female, anterior; 59, tibia and tarsus of palp, female, posterior; 60, chelicera, female; 61, entire palp, female, anterior.

FIGS. 62-64. *Centrotrombidium hadroseta* n. sp., female: 62, ocular plates, anterior aspect on left, dorsal aspect on right; 63, propodosoma; 64, hysterosomal setae, side and top view.

somewhat shorter. Sensilla with terminal portion spherical, rather abruptly differentiated from stalk as in *C. distans*. Ocular plates (Fig. 62) with a single hyaline cornea plus a large punctate posterior lobe bearing a few tubercles. Dorsal body setae borne on individual sclerites as in other members of the genus, the sclerites in the form of very low cylinders, relatively lower than those of *C. distans*; shaft of setae short, extending only a little beyond the margin of the setigerous sclerite, appreciably thicker than in *C. distans*, and not

tapering uniformly as in that species (hence the name). Other details of dorsum and venter essentially as in *C. distans*. Genital sclerites with 7 and 8 setae each, paragenital sclerites with 15 and 16 setae each. Anal sclerites each with 7 setae. In the holotype female, the solenidion of the tarsus of the palp is located somewhat more distally than in the previously described species, and the terminal eupathidia of the tarsus are somewhat longer; however, it is impossible at present to know whether these differences are of real significance.

The material available is not suitable for counts of the setal types. In general the chaetotaxy appears to resemble that of *C. distans* very closely. Tarsus I bears either 2 or 3 clavate solenidia₂, with the normal two being located at 0.67 and 0.74 p in the specimen studied. One of the tarsi I of the holotype has a third s_3 at 0.52 pd . Eupathidia of tarsi I extending from 0.34 v to 0.83 pd . Famulus elongate, straight, its alveolus at 0.88 pd . Claw fossa extending from 0.72 d to end of tarsus. Tarsus II with a clavate s_2 at 0.61 pd and a spikelike famulus at 0.81 pd . These positions correspond almost precisely with the positions of the same setae in the holotype of *C. distans*, but the famulus appears to have a normally recessed alveolus, and not a vesicular one as in *C. distans*.

TYPE LOCALITY: About 4 miles north of Tonasket, Washington, on border of a small permanent pond (holotype female). August 14, 1952, collected by the writer. This is also the type locality of *C. approximatum* new species. Type in author's collection.

REMARKS: The type locality of this species is the only point at which more than one species of *Centrotrombidium* has been observed by the writer, all three forms newly described in this paper having been found here. The differences between *C. distans* and *C. hadroseta* cannot be fully evaluated at present because of lack of sufficient specimens of the latter species. However, 50 specimens of both sexes of *C. distans* have been seen by the writer, and there is no indication of convergence in setal type between any of these specimens and the type of *C. hadroseta*. While it is possible that further study will change this, the most probable explanation at present appears to be that we have here two sibling species.

Centrotrombidium misellum

(Berlese) 1918, new combination

This was originally described as *Diplothrombidium misellum* Berl. (*sic*). A translation of Berlese's description is given above.

395. *Diplothrombidium misellum* Berl. n. sp. Dark red, small. Abdomen subquadrate-rounded, very slightly excavated in lateral margin, front truncate, well-armed posteriorly. Crista metopica with a single area sensilligera, anterior and posterior sensilla very close together; anterior sensilla spiniform, moderately swollen, posterior sensilla long, very slender. Body completely clothed with hemispherical papillae, setae short arcuate, directed posteriorly. Legs very short, anterior ones shorter than body, tarsus broad, ovate, little longer than broad (140 μ long, 80 μ broad), tibia three times narrower than the tarsus. Palpi rather large, barely swollen, unguiculate, provided with a stout spine at the base of the main claw three times shorter than the claw. Papilla, (tarsus?) conical, short, apex armed with three spiniform setae, 30 μ long, cuticle of legs, rough, warty, covered with simple short setae. Eggs in hysterostoma of the single specimen which I saw spherical, 140 μ in diameter. Adult 800 μ long, 650 μ wide.

HABITAT: A single specimen collected by Cl. Alph. Dugès, in Mexico (Guanajuato), which Cl. Trouessart kindly sent to me.

The swollen form of the leg tarsi, the presence of a single pair of true sensilla on the scutum, and the description of the tarsus of the palp and its terminal setae leave little doubt that this is not a *Diplothrombidium* but more likely a *Centrotrombidium*. The eupathidia at the end of the tarsus of *C. approximatum* are almost exactly 30 μ long, while those of *Diplothrombidium micidium* are 62 μ long. No known species of *Diplothrombidium* has a palpal tarsus which could be described as "conical," but this term could be applied to *Centrotrombidium*. The length of the body is also much more in keeping with *Centrotrombidium* than with *Diplothrombidium*. The species should therefore be transferred to the genus *Centrotrombidium*. If it does not belong in that genus, its position there is at least no more untenable than it was in *Diplothrombidium*. The only point in which Berlese's diagnosis was out of character with *Centrotrombidium* is in the form of the sensilla. However, a trend toward more slender sensilla is seen in *C. australasiae*, and it is not inconceivable that there might be some species in which these are not swollen.

Johnstoniana George 1909

ADULT: Large mites, idiosoma 1,000 to 3,000 μ in length, scutum a broad, roughly pentagonal plate with a prominent anterior spine, two pairs of slender sensilla and a fairly well-developed crista metopica. Lateral to the posterior sensilla is a group of several smooth setae, and in the posterior half of the plate usually one or two somewhat isolated setae. Ocular plates protruding, bicornate; propodosoma lateral to scutum devoid of setae. Dorsal hysterosomal setae simple, smooth, borne on individual sclerites which may be elevated or not. Coxae I without supracoxal setae; intercoxal area with numerous smooth setae borne on individual sclerites; pars medialis coxae present. Genital and paragenital sclerites well developed, pregenital tubercle absent, three pairs of genital acetabula. Anal sclerites absent in *J. latiscuta* new species. Only two pairs of rostral setae, the deutorostrals apparently absent. Trochanter of palp not fenestrated, considerably longer anteriorly than posteriorly. Palpal tarsus with a single solenidion on posterior surface, and a number of prominent eupathidia. Palpal tibia with a simple terminal spiniform seta, and a single subterminal spiniform seta. Solenidial types s_1 and s_2 distinct, but s_1 and s_4 difficult to differentiate; s_2 relatively short, stout, marked with four longitudinal ridges. Eupathidia present on all segments of all legs including and beyond the telofemur. Vestigial setae present on patella I and II, but absent on tibia I and II.


LARVA: Scutum bearing two pairs of sensilla plus two pairs of other setae, all simple, smooth. Crista metopica present; anterior end of scutum drawn out into a frontal spine. Supracoxal setae absent. Urpore present. Tip of rostrum with only two pairs of setae, the deutorostrals apparently absent. Tarsus of palp with a single solenidion on posterior surface; typical eupathids absent, although the terminal setae of the tarsus are probably modified eupathids. Femora I to III incom-

pletely divided dorsally in *J. latiscuta*. Patella I and II each with a vestigial seta dorsally, but tibia lacking vestigial setae. Dorsal eupathid of tarsus I with a basal companion seta. Tarsi of all legs with two unequal claws.

REMARKS: The structure of the adult scutum as delineated by Berlese and reproduced by various authors is incorrect. Because of the very sharp downward flexure of the scutum between the posterior and anterior sensilla, the configuration of the scutum in undissected specimens is noticeably different from that seen in dissected specimens (Figs. 70, 86). Little is known of the biology of our one North American species except that it is apparently a cold stenothermal species found along mountain streams. *J. errans* (Johnston) 1852 has larvae which are parasitic upon Tipulidae. Cooreman (1952, p. 109) reported only one case in which he found this ordinarily rare species in great numbers at one point along a small stream in Belgium. This rarity is apparently characteristic of the species described here, for the writer has seen only one female and a larva of *J. latiscuta*. Cooreman reported the capture of 44 females, but no males, which points up the fact that males have not yet been reported for this genus.

Johnstoniana latiscuta new species

FEMALE: Body of ovigerous female about 1816 μ long to tip of scutum, 1330 μ wide, length/width 1.37; color in life a brilliant brick red (Fig. 65). Scutum (Figs. 70, 86) with two pairs of slender, smooth sensilla; plate divided into two portions, a broad extensive posterior portion bearing the posterior sensilla and lateral and posterior to these six to eight setae on either side. The anterior portion is set off by a sharp declivity and bears only the anterior pair of sensilla, in front of which is a rounded knob produced ventrally into a sharp spine. Crista metopica well developed in anterior half of scutum, feebly developed to absent in posterior half. Ocular plates protruding, prominently bicornate;



FIGS. 65-73. *Johnstoniana latiscuta* n. sp., female: 65, dorsum; 66, gnathosoma; 67, genital area; 68, anal area; 69, palp, anterior; 70, scutum; 71, tip of rostrum, ventral; 72, legs I and II; 73, legs III and IV.

cuticle between ocular plates and scutum devoid of setae. Dorsal and marginal body setae (Fig. 84) borne on prominently elevated setigerous sclerites uniformly distributed over the distinctly striated membranous cuticle.

Coxa I with about 60 simple, smooth setae, plus 6 to 9 simple, smooth setae on the pars medialis coxae which appears to be a discrete medial expansion of I (Fig. 82). Supracoxal seta I absent. Coxa II with 45 setae. Membranous area between coxae I and II of right and left sides with 79 smooth setae, borne on individual sclerites. Behind coxa II a wide (75μ) band of striated cuticle devoid of setae. Marginal setae especially dense between II and III, borne on sclerites which are markedly convex but not cylindrical. In the single specimen available, no trace was found of the characteristic organ just anterior to coxa III in *Lassenia lasseni*. Coxa III with 41 setae; IV with 55 (Fig. 75). No apodemes visible between II and III. Genital opening (Fig. 67) large, with three pairs of genital acetabula; genital sclerites very narrow, crescentic, bearing 34 to 36 slender, smooth setae, mostly in a single to double row. Paragenital sclerites with 52 to 55 similar setae in a single to triple row. The setae on both sets of sclerites are fairly uniformly spaced except at the ends where there are fewer setae than at levels in between. Ventral body setae borne on hemispherical sclerites, membranous cuticle marked with regular but not parallel striae. Anus (Fig. 68) lacking well-defined sclerites, surrounded by cuticle only faintly differentiated from that of rest of venter.

Velum circular to oval in outline, numerous fine filaments converging from a well-defined marginal rim (Fig. 71). Protorostral and tritrostral (? see larva) setae present and well developed; ventral surface of gnathosoma


otherwise with 80 to 100 smooth simple setae on either side of the bare mid-ventral line (Fig. 66). Rostrum very short, blunt, extending scarcely beyond the end of the trochanter of the palp.

Chelicerae (Figs. 74, 85) with distal half of tarsal claw slightly raised, knifelike, appearing minutely serrate under oil immersion. Trochanter of palp (Figs. 66, 69) devoid of setae and also without any trace of the fenestration which characterizes the palpal trochanter of the other genera; trochanter considerably longer anteriorly than posteriorly. Terminal segments of palp covered with many long, slender, smooth setae. Tibia with a unidentate terminal seta and a stout subterminal seta at $0.67v$, the latter bearing a basal tooth or spur. A few of the dorsal setae of the tibia, at least in the distal half, are slightly heavier than the other setae, but are not otherwise different from them. Palpal tarsus with a large eupathid at $0.57d$, and five other eupathids at $0.73a$, $0.81d$, $0.94v$, $0.95p$, and $0.97d$. A single solenidion at $0.84p$ (Fig. 83).

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae, m = many).

All segments of all legs beyond the basifemur with one or more eupathidia, recognizable by their reverse curvature. Vestigial setae fairly long and slender, about half the length of the surrounding solenidia₃, confined to patella I and II, absent from tibiae. Tibiae with numerous s₃ (Figs. 77, 78) I and II with only about one seta each which could be assigned to type four. Tarsus I with three s₁ at $0.73d$ to $0.76pd$, plus four others at 0.27 to $0.33d$ which may possibly belong in this category. Solenidial types s₁ and s₄ are convergent in form and difficult to differentiate. Tarsus I

	tr	bf	tf		pa			ti				ta						
	n	n	s ₃	e	s ₃	e	v	s ₃	s ₄	e	v	s ₁	s ₂	s ₃	s ₄	e	f	
I	18	m	2	1	63	3	1	m	1	4	0	3-7	7	0	m	m	1	
II	33	m	1	1	20	3	1	20	1	2	0	4	1	0	10±	20±	1	
III	30	m	1	1	26	2	0	16	0	3	0	0	0	0	7±	3,4	0	
IV	40	m	1	1	35	2	0	28	0	2	0	0	0	5±	0?	2	0	



FIGS. 74-86. *Johnstoniana latiscuta* n. sp., female: 74, end of chelicera; 75, coxae III and IV; 76, solenidion, tibia I; 77, s₃, patella I; 78, s₃, femur I; 79, s₁, tarsus I; 80, s₁, tarsus II; 81, s₂, tarsus I; 82, coxae I and II; 83, tibia and tarsus of palp, posterior; 84, hysterosomal setae; 85, chelicera; 86, propodosoma.

also with seven s_2 at $0.72pd$ to $0.92pd$, these being short, slightly clavate, and longitudinally ribbed; s_3 absent, s_4 numerous. Many eupathidia extending from $0.36v$ to $0.20d$. Famulus₁ at $0.70d$ very similar to the eupathidia in form, but somewhat shorter and broader at the base. Tarsus II with four solenidia₁, at 0.23 , 0.38 , 0.54 and $0.57d$; a single s_2 at $0.60d$, at the base of the claw fossa. Precise number of s_4 not determinable with material available; eupathidia numbering about 20. Famulus₂ at $0.57d$ very small, short, peglike, easily overlooked. Tarsus III with about seven solenidia intermediate to types s_3 and s_4 , but more closely resembling the latter. Tarsus IV with about five solenidia, which appear to belong to type s_3 .

MALE: Unknown.

LARVA: Only one specimen of the larva was taken and this was in such poor condition that neither measurements nor counts of the numerous body setae could be made. Scutum (Fig. 90) with two pairs of smooth, slender sensilla and two pairs of finely barbed setae (oil immersion). Anterior end of scutum with a knoblike projection bearing at its tip a smaller conical projection. Crista well developed; extending to level of anterior sensilla. Cuticle finely and densely punctate. Dorsal and marginal body setae slender, tapering to a very fine point, and smooth; each seta borne on an individual sclerite which is rather feebly delimited from the surrounding unstriated membranous cuticle.

Coxa I with two setae, including the one on the rather well-developed pars medialis coxae, II and III each with a single seta, that on II bifurcate on both right and left sides of the single specimen available (Figs. 87, 101). Urpore well developed. Supracoxal setae on I apparently absent but material not very


satisfactory for study.

Gnathosoma bearing only two pairs of setae, the protorostrals and probably the tritrostrals, the deutorostrals apparently absent (Fig. 88). Supracoxal setae absent. Rostrum short, blunt; velum simple. Chelicerae short, compact, tarsus scythe-shaped with one or two small teeth near the tip. Palpal trochanter (Fig. 94) without setae, femur and patella with a single seta dorsally which appears smooth at low magnifications, but at high magnifications can be seen to have three or four very minute barbs. Tibia with three normal setae in basal half, plus the bidentate terminal clawlike seta. Tarsus with a solenidion very near the base, three long, slender, normal setae also in basal half plus four heavy pectinate setae in distal half. Typical eupathids are absent, although the four hemipectinate setae in the distal half of the segment evidently represent the eupathids of the adult.


Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae).

Femora I to III (Figs. 87, 91, 93) incompletely divided; with a well-developed synarthrodial membrane ventrally but the cuticle in the dorsal part of the segment continues with no trace of interruption from the basifemoral to the telofemoral portion. Although there is no true articulation dorsally as found between other segments of the leg, there is undoubtedly considerable flexibility at this joint. Telofemoral portion of femur, patella and tibia of all legs with one or more solenidia₃ dorsally. Patella I and II each with a vestigial seta dorsally, but tibiae lacking vestigial setae. Tarsus I (Fig. 99) with annulate dorsal solenidion₁ at $0.45d$, a straight, elongate, smooth famulus at $0.56pd$, eupathid and

	tr	bf	tf		pa				ti				ta					
	n	n	s ₃	n	s ₃	s ₄	v	n	s ₃	s ₄	v	n	s ₁	s ₂	e	f	c	n
I	1	1	1	5	4	0?	1	4	0	2	0	6,7	1	0	2	1	1	37
II	1	2	1	4	2	0?	1	4	1	1	0	6	0	1	1	1	0	28
III	1	2	1	4	2	0	0	4	1	1	0	6	0	0	0	0	0	23



FIGS. 87-98. *Johnstoniana latiscuta* n. sp., larva: 87, leg I, and coxae I and II; 88, gnathosoma, ventrolateral; 89, tibia and tarsus of palp, posterior; 90, scutum; 91, leg II, trochanter to tibia; 92, tip of palpal tarsus; 93, leg III, trochanter to tibia; 94, palp, posterior; 95, solenidion, patella II; 96, s₁, tarsus I; 97, s₂, tarsus II; 98, s₄, tibia I.



FIGS. 99-102. *Johnstoniana latiscuta* n. sp., larva: 99, tarsus I, anterior; 100, tarsus II, anterior; 101, coxa III; 102, tarsus III.

companion seta at $0.70d$, and a shorter one at $0.88pv$. Tarsus II with short, spikelike famulus₂ at $0.50pd$, a clavate, carinate solenidion₂ at $0.59pd$, and a rather straight eupathid at $0.84pv$. Tarsus III with no specialized setae. Anterior and posterior claws on all legs with a sharp spine at the basal third of the concave side; anterior claw gently curved throughout length, posterior claw sharply bent just beyond the spine.

TYPE LOCALITY: At origin of King's Creek, Lassen Volcanic Park, California, August 6, 1955 (holotype female). Collected by the writer. Type in author's collection.

REMARKS: The correlation of the female with the larva described above must be regarded as tentative, although it is almost certain that they belong to the same species. The two specimens were found within about a foot of each other under rocks on the edge of the stream. There is no question about their generic identity.

It is not possible to compare this species with other presently described forms because

of the inadequacies of the description of the latter. However, it is probable that detailed comparisons would prove it distinct.

Johnstoniana vitzthumi Womersley 1939

While it is apparent from the original figures and description that this species belongs in the Johnstonianidae, it is impossible to compare it reliably with other members of the family, or even to ascertain that it is truly a *Johnstoniana*. The description was based on a single specimen collected in South Australia.

Diplothrombium Berlese 1910

ADULT: Fairly large mites, idiosoma usually between 1,000 and 2,000 μ long, color in life dark reddish-brown. Scutum elongate, narrow, bearing an anterior spine, and well-differentiated anterior and posterior areas sensilligeræ (Figs. 110, 160). Two pairs of smooth, slender sensilla, and usually two pairs of shorter normal setae in the posterior area sensilligera. Dorsum of propodosoma with a few to many setae borne on individual

sclerites between the scutum and ocular plates. Dorsal hysterosomal setae smooth, borne on individual elevated sclerites. Ocular plates protruding moderately, bicornate. Coxa I lacking supracoxal setae; coxal ring of II complete dorsally, that of coxa I membranous dorsally. Intercoxal area with numerous smooth, slender setae, each borne on individual sclerites; pars medialis coxae well developed. Genital and paragenital sclerites well developed, three pairs of genital acetabula. Anal sclerites crescentic, bearing a number of smooth setae. Rostrum apparently only with the protorostral and tritrostral setae, plus other setae behind these; deutorostrals absent. Trochanter of palp fenestrated anteriorly; tibia with one large terminal and one subterminal clawlike seta. Tarsus of palp with a single solenidion posteriorly, and a number of eupathidia in the distal half. Legs with solenidia of four fairly distinct types, s_2 being especially distinctive, clavate. Eupathidia present on all segments of all legs including and beyond the telofemur. Vestigial setae completely absent.


LARVA: Scutum with two pairs of sensilla, the anterior pair very small; otherwise with two pairs of normal setae. Crista metopica well developed, anterior portion of scutum produced into a process of variable form. Ocular plates bicornate. Dorsal and ventral surfaces of body covered with numerous setigerous sclerites. Coxal setae numbering 2–1–1 in known species. Intercoxal area with a single pair of setae between III; urpore present. Gnathosoma bearing only two pairs of setae, probably the protorostrals and tritrostrals. Supracoxal setae absent on both the gnathosoma and on leg I. Podocephalic canals well developed. Anterior wall of trochanter of palp fenestrated. Palpal tarsus with a single solenidion near basal end; eupathidiform setae absent, although some of the terminal setae must be modified eupathidia. Solenidia₃ present on telofemur of legs I to III; s_1 elongate, somewhat decumbent, s_2 stout, clavate. Tarsus II with only one eupathid. All

legs with six free segments beyond the coxa, and two subequal tarsal claws.

REMARKS: The larvae described here are the first described for the genus as a whole, and, of course, *D. monoense* is the first species in which the larva and adult are both known. Species of this genus are found in very moist situations and are cryptic in habits, avoiding direct sunlight as much as possible. Although experimental evidence is not yet available, they are evidently incapable of tolerating high temperatures for any length of time. The larvae are of the self-detaching type; they have been found parasitizing the pupae of aquatic beetles under wet rocks.

Diplothrombium monoense n. sp.

FEMALE: Idiosoma (Fig. 103) highly variable in size, 1,638 to 1,820 μ long, 1,092 to 1,326 μ wide, length/width 1.43 to 1.52; average 1,706 by 1,180 μ , length/width 1.45 (five ovigerous specimens). Scutum (Fig. 110) with two pairs of completely smooth sensilla, and between them a pair of stiff setae appearing hemipectinate under high magnification. Posterolateral to these is a second pair of setae contiguous with, but not fully enclosed by the scutum. Anterior end of scutum produced into a long blunt spine. Crista metopica well developed, extending from posterior margin of anterior area sensilligera to posterior end of scutum. Behind the posterior area sensilligera is a lateral expansion on either side of the crista metopica, bearing a number of coarse punctations. Ocular plates small, scarcely larger than the two corneae found on each side, strongly convex but not stalked, setae absent. Dorsal and marginal body setae (Fig. 109) smooth, slender, tapering, typically with a sharp basal flexure. Alveoli borne on individual sclerites which are sharply elevated above the general surface of the cuticle in the form of a truncate cone. Propodosomal cuticle between the scutum and ocular plates and lateral to the ocular plates generally with



FIGS. 103-112. *Diplothrombium monoense* n. sp.: 103, dorsum, female; 104, venter, male; 105, anus, male; 106, gnathosoma, male, lateral; 107a, coxae III and IV, male, lateral; 107b, coxae I and II, male, lateral; 108, chelicera, female; 109, hysterosomal seta, female, side view; 110, propodosoma, female; 111, genital opening, female; 112, genital opening, male.

fewer than ten setae. Cuticle seemingly devoid of striae or other markings except for the underlying reticular layer.

Coxae I and II (Fig. 122) with about 50 and 60 setae respectively, plus 7 to 10 setae on the pars medialis. All setae smooth. No supracoxal setae; coxal ring I open dorsally, II closed dorsally (Fig. 107b). Coxae III and IV with 70 to 75 setae each, coxal ring of each complete dorsally (Fig. 107a). Cuticle of all coxae faintly punctate. Intercoxal area between I and II with 65 to 70 smooth slender setae, each borne on a small sclerite; a narrow band of cuticle behind coxae I and II devoid of setae. No sclerotized plates in the membranous area between coxae II and III. Ventral body setae like those on dorsum, except that the sclerites are not so prominently raised. Genital sclerites (Fig. 111) bearing 35 to 45 smooth slender setae; paragenital sclerites narrower than genital sclerites but bearing 43 to 50 setae of the same form as those on the genital sclerites (three specimens). Genital acetabula well developed, the second pair the smallest. Anal sclerites (Fig. 105) quite variable, with from 11 to 19 setae arranged in a single to double row; setae similar to those on genital sclerites.

Base of gnathosoma and rostrum bearing about 45 setae on each side, supracoxal setae absent. Rostrum relatively short and broad; tip of rostrum (Fig. 106) with two pairs of specialized setae, presumably the protorostrals and tritrostrals. Velum small, circular, and oriented in an anterior direction (Fig. 125). Chelicerae as shown in Figure 108. Trochanter of palp (Fig. 127) devoid of setae; medial wall fenestrated. Tibia with a heavy spiniform seta at 0.58*v* and a second one at the end of the segment. A series of three very sharp spines dorsally, in basal third of tibia


(Fig. 126). Tarsus with a large eupathid borne on a prominent tubercle at 0.61*d* and a group of two large and two small eupathids near the tip of the segment (Fig. 124). A single solenidion at 0.57*p*. Normal setae of palp smooth and tapering.

Trochanters I and II with about 8 setae dorsally, three with about 10 setae dorsally, and four with 20 to 25 setae, all these setae smooth to faintly hemipectinate. All basifemora lacking specialized setae, normal setae on dorsal portions of segment smooth to faintly hemipectinate, those on ventral half of segment usually very fine and smooth.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae, m = many).

Vestigial setae absent from all legs. Tarsus I with famulus elongate, erect, its alveolus located on a prominent tubercle at 0.31*pd* (Fig. 123). Solenidion₁ actually smaller than the corresponding seta on tarsus I of the larva, inserted at 0.52*d* at the origin of the claw fossa. A group of five thick, stout *s*₂ extending from 0.57 to 0.80*p*. These are generally slightly smaller than the corresponding setae on tarsus II but are the same general form and are internally annulate. Eupathidia very numerous, extending from 0.26*v* to 0.13*d*, or in other words around nearly the entire periphery of the tarsus I. Basal portion of tarsus I from about 0.30*v* to 0.52*d*, extremely rough, provided with sharp, truncate tubercles. These tubercles bear the alveoli of a number of types of setae including the famulus, eupathidia, solenidia₄, and bihemipectinate normal setae. The normal setae on all segments of the legs are smooth or very nearly so, with the exception of those of tarsus I. Tarsus II with a clavate *s*₂ at 0.49 to 0.50*d*, and a spikelike

	tr	bf	tf		pa			ti			ta					
	n	n	s ₃	e	s ₃	s ₄	e	s ₃	s ₄	e	s ₁	s ₂	s ₃	s ₄	e	f
I	7-9	m	16	3	60	2	2	55	2	2	1	5	0	m	m	1
II	7	m	10	3	22	3	3	10	5	3	0	1	0	5	4	1
III	9-11	m	10	3	30	1	2	11	5	3	0	0	0	2	8	0
IV	20-23	m	6	2	30	2	3	13	4	4	0	0	0	4	8	0



FIGS. 113-128. *Diplotbrombium monoense* n. sp.: 113, legs I and II, female; 114, legs III and IV; 115, solenidion₃, telofemur I; 116, s_3 , tibia I; 117, s_4 , tarsus I; 118, s_1 , tarsus I; 119, s_2 , tarsus II; 120, s_2 , tarsus I; 121, coxae III and IV, female (115-120 all female); 122, coxae I and II, female; 123, famulus and other setae of tarsus I, female; 124, tibia and tarsus of palp, female, posterior; 125, gnathosoma, female; 126, tibia of palp, female; 127, entire palp, female; 128, tarsus II, female.

famulus at 0.45 to 0.52*d*, these setae of the same form as the corresponding ones in the larva. Tarsus II apparently lacking *s*₁, the remaining solenidia being *s*₄ (Fig. 128). All tarsi with a distinct claw fossa on dorsal surface, and most segments of legs with exception of trochanter and basifemur bearing many angular dorsal protuberances which give the legs a rough appearance. Two claws on all tarsi, the claws flattened and with a median ventral carina.

MALE: Similar to female in most respects observed, with the principal exception of the genital opening (Fig. 112). Genital opening guarded by two pairs of sclerites as in female, but these have many more setae. Anterior 0.4 of each genital sclerite with 45 to 50 close-packed smooth, long, slender setae; remaining 0.6 with only about 20. Paragenital sclerites each with about 40 to 45 smooth setae, 6 or 8 of which are appreciably heavier than the others. Penis not triangular, but small and tubular. Three pairs of prominent genital acetubula. Anal sclerites with 14 to 17 smooth, slender setae.

LARVA: Body 343 to 389 μ long, 233 to 279 μ wide, length/width = 1.36–1.57; average 369 by 253 μ , length/width = 1.46 (five specimens). Scutum (Fig. 129) with a knob-like protuberance anteriorly, and a pair of very small sensilla inserted near the base of this. The knob bears a short, blunt point ventrally, but this is not visible in dorsal view. These setae, while not especially sensillar in form, are forerunners of the anterior sensilla in the adult. Setae at anterolateral and posterolateral angles bihemipectinate. Posterior sensilla very long, slender, smooth. Crista metopica well developed in anterior one-half (sometimes in posterior one-third also) of scutum. A patch of moderately coarse pores (250 x) near posterior margin; surface otherwise marked with numerous fine punctae and an irregular reticulum medially behind the sensilla. These pores and reticulae are also evident in the adult (Fig. 110). Corneae two on each side, borne on a common ocular plate. Dorsal setae

36 in number, borne in five transverse rows (8, 8, 8, 8, 4; Fig. 131). Each seta faintly hemipectinate (250 x), borne on an individual sclerite, the central portion of which is moderately elevated (Fig. 130). Membranous cuticle with parallel striae, which are sometimes indistinct.

Coxa I (Fig. 144) with two bifurcate setae; medial portion of coxa caudiform, indistinct. Supracoxal seta absent. Coxa II with one bipectinate seta laterally; coxa III with one smooth seta. Intercoxal area (Fig. 138) with only one pair of setae between coxae III; postcoxal area with 36 to 38 hemipectinate ventral and marginal setae on each side, all borne on individual sclerites. Presumptive anal opening distinct.

Gnathosoma with a single pair of bifurcate setae ventrally, presumably the tritorostrals. Rostrum very short; protorostral setae smooth. Velum small. Podocephalic canals well developed. Supracoxal setae absent. Cheliceral base very thick; digitus fixus membranous (Fig. 134); digitus mobilis smooth, except for one tooth. The ventral margin of the digitus mobilis in the specimen drawn was bent, but this may not be normal. Palpi five-segmented (Fig. 133). Trochanter lacking setae, femur and patella each with one long, smooth dorsal seta. Tibia with three smooth setae and a hooked terminal spiniform seta which usually appears unidentate in lateral view, but bifid in ventral view. Palpal tarsus (Figs. 132, 136) with solenidion basally on lateral surface. Tip of tarsus drawn out into a flat, sharp spine; a faintly pectinate seta arising dorsally at the base of the spine gives the tarsus a bifid appearance. Otherwise with six setae, all but two of which are faintly hemipectinate. No typical eupathidiform setae present, although the heavy pectinate setae are quite certainly the forerunners of these.

Chaetotaxy of legs approximately as shown in table (*s* = solenidia, *e* = eupathidia, *f* = famulus, *n* = normal setae).

Trochanters I to III each with one large ventrally curved seta. Basifemora all distinctly

	tr	bf	tf		pa		ti			ta				
	n	n	s ₃	n	s ₃	n	s ₃	s ₄	n	s ₁	s ₂	e	f	n
I	1	1	2	5	15	4	1	1	6	1	0	2	1	34
II	1	2	1	4	2	4	0	2	6	0	1	1	1	27
III	1	2	1	4	2	4	0	1	6	0	0	0	0	20


separated from telofemora; I with a large ventral seta, but no dorsal seta, II and III each with a large hemipectinate dorsal and a small, smooth ventral seta. Telfemur I with two dorsal solenidia₃ (Figs. 139–141), II and III with one each; otherwise with 5, 4, and 4 smooth to faintly pectinate setae respectively. Vestigial setae and companion setae absent from all segments of all legs. Tarsus I (Fig. 143) with a long famulus at 0.42 pd and a long solenidion₁ at 0.44 d , the latter lacking annular structure, but with internal surface granular (oil immersion). Solenidion, when seen in dorsal view, not straight, but very gently bisinuate. A long eupathid at 0.66 d , but no companion seta, and a shorter one at 0.93 v . A pair of long, slender setae at 0.87 and 0.89 ad and pd , and another at 0.88 av . Otherwise with 31 bihemipectinate setae in specimen drawn. Tarsus II (Fig. 145) with famulus at 0.45–0.52, a club-shaped solenidion₂ at 0.55 d , three slender, smooth setae at 0.80 d , 0.83 d , and 0.83 v , and a eupathid at 0.89 pv . Otherwise with 24 bihemipectinate normal setae in specimen drawn. Tarsus III (Fig. 142) with five slender, smooth setae at 0.79 d to 0.86 v . Otherwise with 15 bihemipectinate normal setae on tarsus drawn. Two claws on all tarsi, the anterior one larger than the posterior one; all claws smooth.

TYPE LOCALITY: Mono Lake, Mono County, California, 0.9 miles north of Leevining (holotype female). Type in author's collection.


REMARKS: These mites were first found on and under rocks, lying in and near a small tricklet of alkaline water running into the lake, June 1, 1952. The habitat could be termed semiaquatic. The bottom of one rock was covered with scores of pink eggs laid singly and several other rocks were also observed with smaller numbers of eggs but

always in association with these adults (June 1, 1952). Eight larvae were found attached to the pupa of an aquatic beetle found under a stone. The larvae were definitely attached to the host and appeared to be feeding, but when submerged in alcohol the larvae immediately detached themselves from the host. The feeding habits of the adult are unknown at present. The same spot was revisited July 1, 1954. A few adults were found at this time, but no eggs, indicating that the reproductive season terminated sometime during the month of June. The same picture was observed when the spot was again revisited August 9, 1955 —although adults and nymphs were fairly common, no eggs or larvae whatever were found.

While found at a site in which very high summer temperatures are registered in exposed places, *Diplothrombium monoense* was never found out in the open where it would be subject to these high temperatures. Also it was never found in situations where there was not considerable moisture, and the majority were found under rocks in which the surface of the ground was actually wet. A few individuals were running about over the surface of very wet ground covered with a dense growth of small plants. Under such circumstances they always appeared restless, especially when their cover was removed so as to expose them to the direct rays of the sun. A species of *Microtrombidium* was also very common at this locality but was frequently found running about over the surface of the dry sand along the shore of the lake, even in the heat of the day. Their movements were generally rapid, but at least they did tolerate the exposure to heat and sun for short periods of time. This species of *Diplothrombium* would appear to be considerably less tolerant of heat and dryness than species of *Microtrombidium*. Fur-



FIGS. 129–138. *Diplothrombium monoense* n. sp., larva: 129, scutum; 130, dorsal seta; 131, dorsum; 132, tibia and tarsus of palp, posterior; 133, entire palp, posterior; 134, end of chelicera; 135, gnathosoma, ventral; 136, tibia and tarsus of palp, anterior; 137, ocular plate; 138, venter.



FIGS. 139-151. *Diplothrombium monoense* n. sp., larva: 139, telofemur to tibia I; 140, telofemur to tibia II; 141, telofemur to tibia III; 142, tarsus III; 143, tarsus I, posterior; 144, coxae I and II; 145, tarsus II, posterior; 146, tip of tarsus; 147, solenidion₂ tarsus II; 148, s₁ tarsus I; 149, s₂ femur II; 150, s₄ tibia I; 151, s₄ tibia I.

ther evidence of the narrow temperature tolerance of *D. monoense* was obtained through an unfortunate mishap. A number of adults being brought back to Riverside for rearing studies were killed during the eight-hour transit of the Mojave Desert, despite precautions taken to insulate them from the heat. The same was true of the living specimens of *Lassenia lassenii*—none survived the trip across the desert. At the same time a species of *Microtrombidium* collected at an elevation of 8,000 feet on the slopes of Mt. Lassen did survive, laid eggs, and these eggs eventually hatched. A species of Erythraeidae also survived the exposure to high temperature which was lethal to the two species of Johnstonianidae. The rather low heat tolerance of these species undoubtedly is important in determining their distribution, both geographically and locally.

The correlation between larva and adult was established on the basis of their co-existence at the type locality. More than 90 adults or nymphs and 50 larvae have been collected there, and there is apparently only one species involved.

Diplothrombium micidium new species

FEMALE: Body (Fig. 152) 936 μ to tip of scutal spine, 546 μ wide, length/width 1.48 (one ovigerous female). Scutum (Fig. 160) produced into a stout conical spine anteriorly. Crista metopica well developed, extending from posterior end of scutum to base of spine. Two pairs of sensilla present, the posterior pair borne on the widest portion of the scutum; sensilla completely smooth. Just in front of the posterior sensilla are two pairs of smooth setae, and just lateral to the sensilla a third pair of setae which lie either in the very margin of the scutum or possibly in some cases just outside. The rather broad plate found behind the posterior area sensilligera in *Diplothrombium monoense* is also found in this species but it is small and easily overlooked at low magnification. Ocular plates

very small, bearing two protruding corneae; devoid of setae. Dorsal propodosomal cuticle containing about 20 setae on each side between the ocular plate and the scutum; each of these setae is borne on a small sclerite. Dorsal hysterosomal setae (Fig. 167) also borne on individual sclerites, the shaft sharply deflexed at the origin of the seta from the alveolus. Membranous cuticle of dorsum smooth, subcuticular reticulum present but not very prominent. Coxae I and II (Fig. 154) each with about 30 or 31 smooth simple setae, pars medialis coxae with three to four setae each borne in an oval area within the pars. Outside of these oval areas the pars medialis is distinctly reticular in appearance. Supracoxal setae absent on I. Intercoxal area with about 60 setae borne on individual sclerites in the specimen studied (not all shown in the figure), these setae smooth and simple. Two to three narrow refractile chitinous rods in the membranous cuticle behind coxa II; a band of membranous cuticle behind coxa II devoid of setae. Coxae III and IV (Fig. 165) each with about 25 to 30 setae. Genital sclerites (Fig. 164) with 10 and 12 setae each in a single row; paragenital sclerites with 19 and 22 setae each in a single to double row. Three pairs of genital acetabula present. Anal sclerites narrow, crescentic, bearing 9 smooth setae each in the specimen studied (Fig. 158). Ventral setae of hysterosoma all borne on individual sclerites as on dorsum.

Rostrum (Fig. 154) with telorostral and deutorostral setae well developed; remainder of rostrum and base of gnathosoma with 30 setae on each side of the mid-line; mid-ventral portion of gnathosoma devoid of setae. Velum small, anteriorly directed, with numerous converging filaments. Base of chelicera (Fig. 166) compact, densely and minutely punctate; dorsal membrane blunt, tarsus stout, curved, dorsal margin appearing smooth at low magnifications but with a very large number of minute teeth at magnifications above 250 x. Palpi (Fig. 153) with trochanter fenestrated on anterior surface, femur with 20

to 25 setae, patella with about 20 setae. Tibia of palp (Fig. 159) with a few sharp spines along the dorsal margin, 10 simple, smooth setae, plus the terminal clawlike seta and the heavy subterminal seta on the anterior aspect. Tarsus of palp with a solenidion at 0.50*p*, a eupathid at 0.64*d* and four additional eupathids at or near the end of the segment, plus 27 smooth setae. On the right palp of the specimen drawn there was a sharp spine at 0.79*d*, just anterior to the dorsal eupathid; on the left palp no such spine was found in this position, but one was found at 0.8*v*. Chaetotaxy of legs essentially as described for male.

MALE: Body 1,014–1,066 μ long, 520–598 μ wide, length/width 1.78–1.95 (average of 3 specimens 1,040 by 554 μ , average length/width, 1.88). Resembling female in all essential respects except structure of genital area. Genital sclerites (Fig. 155) with 16 to 18 setae in a single to double row, paragenital sclerites with 23–26 setae, also in a single to double row; an internal cirlet of about 7 pairs of setae. Genital acetabula numbering three pairs. Anal sclerites as described for female.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, n = normal setae, m = many).

All legs with telofemur to tarsus very rough in appearance (Figs. 162, 163). Solenidia exceptionally well defined, with no intergradation between types; *s*₃ slender, with no internal structure, *s*₄ bacilliform with usually distinct internal structure, somewhat variable in length. Solenidia₁ and *s*₂ are not unlike *s*₃ in general appearance, but differ principally in length and moreover occupy characteristic positions on the tarsi. Solenidia₄ absent from all but tarsus of leg I, but present on telofemur to tarsus of legs II to IV. Vestigial

setae absent from all legs. Tarsus I with famulus at 0.65*d*, this seta resembling a eupathid but somewhat shorter. Solenidion₁ at 0.80*d*, and three *s*₂ at 0.63, 0.76, and 0.87*p* (Fig. 161). Tarsus II (Fig. 156) with spikelike famulus at 0.59*d*, *s*₂ at 0.60*d*, and a group of five *s*₄ at 0.18*d* to 0.49*p*. Tarsus II with only two eupathidia, one at 0.54*pd*, the other at about 0.95*pv*. Eupathidia present on all segments of all legs beyond the basifemur, but few in number, rarely exceeding three except on tarsus I. Eupathidia of tarsus I extending from 0.35*v* to 0.39*d*. All tarsi with two slightly subequal claws and a rather small claw fossa.


LARVA: Unknown.

TYPE LOCALITY: East fork of Hat Creek, above Hat Lake, at an elevation of about 7,250 ft., Mount Lassen, California (holotype male), under rocks and sticks along the stream in a cascade meadow. August 7, 1955, collected by the writer. Type in author's collection.

The writer wishes to express his appreciation to the National Park Service for their cooperation in these studies. As a general rule, most forms which can be collected within National Park boundaries can also be collected outside with nearly equal facility. However, three species of great interest in the present work (*D. micidium*, *Johnstoniana laticuta*, and *Lassenia lasseni*) have been found only within the limits of Lassen Volcanic Park, despite intensive collecting at a large number of points in the Pacific coast states.

REMARKS: This species can be readily differentiated from *D. monoense* n. sp. by several characteristics. It is much smaller, the length being about 1,040 μ compared with 1,630 to 1,820 μ for *D. monoense*. There are about 20 dorsal propodosomal setae outside of the scutum, compared with fewer than 10 in most

	tr	bf	tf			pa			ti			ta					
	n	n	s ₃	s ₄	e	s ₃	s ₄	e	s ₃	s ₄	e	s ₁	s ₂	s ₃	s ₄	e	f
I	6	m	19	0	1	53	0	2	58	0	3	1	3	0	41	m	1
II	4,5	m	4	1	1	10	2	1	8	3	2	0	1	0	5	2	1
III	6	m	7	1	1	13	4	2	10	3	3	0	0	0	5	2	0
IV	14	m	7	3	1	17	6	3	16	5	3	0	0	0	5,6	2,3	0



FIGS. 152–161. *Diplothrombium micidium* n. sp.: 152, dorsum, female; 153, palp, anterior; 154, gnathosoma, coxae I and II, intercoxal area, female; 155, genital area, female; 156, tarsus II showing setal types, female; 157, famulus and nearby eupathid, male; 158, anus, female; 159, tibia and tarsus of palp, female, anterior; 160, propodosoma, female; 161, tarsus I, female, posterior.

specimens of *D. monoense*. The genital and paragenital sclerites of the female have considerably fewer setae than in the comparison species; also the genital sclerites of the male have 16 to 18 uniformly spaced setae, while in *D. monoense* there is a dense concentration of setae in the anterior 0.4 of the plate. Famulus₁ is at 0.65*d* while in *D. monoense* it is at 0.29–0.32*d*. The solenidion₁ of tarsus I is at 0.80*d* compared with 0.52*d* for *D. monoense*. Other differences can be found in the descriptions of the two species, or by comparison of specimens.

Diplothrombium cascadenae new species

LARVA: Idiosoma (Fig. 168) 261 to 315 μ long, 189 to 225 μ wide, length/width = 1.38–1.44; average of eight specimens 288 μ by 207 μ , length/width = 1.41. Scutum (Fig. 174) with two pairs of sensilla, the posterior ones long, smooth, the anterior ones very short; otherwise with four faintly pectinate setae. A knoblike projection at the anterior end of the scutum, acuminate in normal specimens, but the point usually directed ventrally and ordinarily visible only in lateral view. This point is much better developed than in *D. monoense*. A distinct transverse line crosses the scutum behind the posterior sensilla. Crista metopica distinct. Corneae two on each side, borne on a feebly developed and protruding ocular plate. Dorsal and marginal setae about 30 in number, each borne on a platelet, shaft smooth, or with a very few fine barbs. Membranous cuticle striate.

Coxal and intercoxal chaetotaxy (Fig. 169) as in *D. monoense*. No supracoxal seta on I. Postcoxal area with only about 20 smooth setae on each side, each borne on a separate plate. Anal anlage well developed, but without sclerites.

Base of gnathosoma as described for *D. monoense*; supracoxal setae absent. Palpal trochanter fenestrated anteriorly, lacking setae (Fig. 179). Femur, patella and tibia with one, one and three slender setae respectively, tibia with large, clawlike terminal seta bifid. Palpal tarsus essentially identical with that of *D. monoense*, but smaller. All setae hemipectinate (Fig. 176). Podocephalic canals well developed.

Chaetotaxy of legs as shown in accompanying table (s = solenidia, e = eupathidia, f = famulus, n = normal setae).

Patella I with only 8 solenidia₃ dorsally, compared with the 15 in *D. monoense*. Tarsus I (Fig. 178) with large, procumbent solenidion₁ at 0.35*d*, eupathidia at 0.59*d*, and 0.89*v*, famulus displaced far anteriorly to 0.65 (the famulus in *D. monoense* is at the same level as the solenidion). Dorsal eupathid without companion seta. A group of three smooth, slender setae at 0.76, 0.80 and 0.84, as in *D. monoense*; otherwise with 29 bihemipectinate normal setae in specimen drawn. Tarsus II (Fig. 175) with short, spikelike famulus at 0.44, a solenidion at 0.47, and a eupathid at 0.86*v*. A pair of smooth, slender setae at 0.77 and 0.80; otherwise with 21 bihemipectinate normal setae.


Tarsus III (Fig. 177) with a group of five smooth, slender setae lying between 0.73*d* and 0.84*v*; otherwise with 11 bihemipectinate setae. All tarsi with two smooth, unequal, scythe-shaped claws.

TYPE LOCALITY: Salt Creek Falls, Lane Co., Oregon. Grass and flowers (holotype larva). July 7, 1952. Collected by the writer. Type in author's collection.

REMARKS: The larva of this species differs from that of *D. monoense* in a number of significant respects. It is smaller, ranging in

CHAETOTAXY OF LEGS

	tr	bf	tf		pa		ti			ta				
	n	n	s ₃	n	s ₃	n	s ₃	s ₄	n	s ₁	s ₂	e	f	n
I	1	1	2	5	8	4	1	1	6	1	0	2	1	32
II	1	2	1	4	2	4	1	1	6	0	1	1	1	23
III	1	2	1	4	2	4	1	0	6	0	0	0	0	16




FIGS. 162-167. *Diplothrombium micidium* n. sp.: 162, legs I and II, female; 163, legs III and IV, female; 164, genital area, male; 165, coxae III and IV, female; 166, chelicera, female; 167, hysterosomal setae, female, side and top views.

FIGS. 168-170. *Diplothrombium cascadenae* n. sp., larva: 168, dorsum; 169, venter; 170, gnathosoma, lateral view.

length from 261 to 315 μ (*D. m.*: 343 to 389 μ), the dorsal and ventral body setae behind the scutum and coxae number only about 50 (*D. m.*: 72-74 setae here), patella I has only

8 s_3 dorsally (*D. m.*: about 15 s_3 here), and the famulus of tarsus I is at 0.65, while the solenidion is at 0.35 (*D. m.*: famulus₁ and s_1 both at about 0.43). Other differences can be



FIGS. 171-179. *Diplothrombium cascadense* n. sp., larva: 171, leg III, telofemur to tibia; 172, leg I, telofemur to tibia; 173, leg II, telofemur to tibia; 174, propodosoma; 175, tarsus II, posterior; 176, tibia and tarsus of palp, posterior; 177, tarsus III, posterior (the smooth seta at 0.74 μ is a normal seta); 178, tarsus I, posterior; 179, entire palp, anterior.

found in the descriptions of the two forms.

No adult of this species was found, despite extensive collections made at the type locality.

Diplothrombium longipalpe Berlese 1887

There is little in the original description of this species which can be relied upon to distinguish critically between this form and the two species described here by the writer. The basal prominence of tarsus I appears to be significantly closer to the base of the segment than in *D. monoense*, but a similar prominence is found in approximately the same position in *D. micidium*. It is probable however that direct comparison of *D. micidium* and *D. longipalpe* would reveal reliable specific differences. Berlese regarded the form named just below as a variety of *D. longipalpe*, but it is likely that if the differences Berlese noted were real that there are two distinct species. A complete bibliography of the species was given by Thor and Willmann (1947, p. 223).

Diplothrombium septentrionale

Berlese 1912, new combination

This form was considered a variety of *D. longipalpe*, but it is almost certainly a distinct species. Again, the information available on this species in the literature is totally inadequate to make any critical comparisons with the two new forms described in the present paper.

Diplothrombium misellum Berlese 1918

Berlese provided no figure of this species, but judging from his description (a single area sensilligera, anterior setae spiniform) this is not a *Diplothrombium* but probably a *Centrotrombidium*. Hence, it is provisionally placed in that genus, *q.v.*

LASSENIINAE new subfamily

DIAGNOSIS: Adults with anterior sensilla

greatly reduced or of markedly different form from posterior sensilla. Pregenital tubercle present. Supracoxal setae present on coxa I and gnathosoma in both larva and adult. Larvae with anal sclerites. Deutorostral setae present; terminal seta of palp eupathidiform. Tarsi each with three claws.

REMARKS: To each of the above statements should be added "in known forms." The group will have to be redefined as it becomes better known.

Lassenia new genus

ADULTS: Rather large mites, the known species between 1,000 and 1,700 μ in length, dark brown to brownish red. Scutum large, expanded, several times wider than the width of the crista metopica. Posterior sensilla elongate, slender; anterior sensilla variable in form, resembling the rest of the setae on the scutum so that they are sometimes not readily discernible. Ocular plates bicornate, protruding, but not stalked. Dorsal setae of hysterosoma simple, smooth, stiff, rodlike in the known species, all of same type. Pars medialis of coxa I either present or absent. Coxal rings I and II both membranous dorsally, coxa I with supracoxal seta. Genital opening with three pairs of acetabula, guarded by a pair of crescentic genital sclerites and a pair of paragenital sclerites; a hemispherical or pedunculate sclerite immediately anterior to the genital opening. Anal sclerites present. A characteristic *Lassenia*-organ anterior to coxa III. Base of gnathosoma with or without setae; supracoxal setae present. Proto-, deuto-, and tritrostral setae all present. Velum simple. Chelicerae slightly to noticeably deflexed posteriorly. Palpi with five free segments, distal margin of trochanter deeply incised on anterior surface. Palpal tibia with heavy terminal seta, at the base of which is a single stout seta. Tarsus of palp with a single solenidion on posterior aspect which may be in either the basal or the distal half of the segment; distal half of the tarsus richly supplied with eupathidia. Eupathidia present on all

segments of all legs beyond the basifemur, sometimes present on ventral surface of telofemur, patella and tibia I. Vestigial setae present on patella I and II and tibia I, but absent on tibia II. Solenidia₁, s₂ and s₄ generally indistinguishable, so that in effect only two solenidial types are present on the legs. All tarsi essentially lacking claw fossa; IV without a posterior eupathid in terminal 1/10th of segment.

LARVA: Parasitic upon Diptera living in subaquatic environments. Scutum of known species with a pair of long posterior, and a pair of shorter anterior sensilla, plus two pairs of other setae near the anterolateral and posterolateral corners of plate; anterior sensilla sometimes set off on a minute sclerite separate from the rest of the scutum. Crista absent. Dorsal setae variable in number, borne on individual setigerous sclerites. Ocular plates bicorneate. Coxae I with two pairs of setae, II with one or two pairs of setae, and III with two or three pairs of setae; a well-developed *Lassenia*-organ anterior to coxa III. Urpore present. Anal sclerites with two pairs of setae in known species. Base of gnathosoma with only the supracoxal setae, rostrum with protorostral, deuterostral and tritrostral setae present and well developed. Palpi somewhat geniculate, femur considerably heavier than rest of segments, and sometimes incompletely separated from patella. Tibia with heavy unidentate or bidentate seta at tip. Tarsus of palp elongate, cylindrical, with a single solenidion posteriorly, a terminal and subterminal eupathid. Femur of legs undivided, legs with only five free segments beyond the coxae. Vestigial setae present on patella I, II, and tibia I, but absent from tibia II. Tibia I with two types of solenidia, s₃ and s₄, the latter larger than the former, with distinct internal structure, and a basal companion seta. Solenidion₁ also with a basal companion seta, s₂ without a companion seta. Dorsal eupathid of tarsi I and II with a basal companion seta. All tarsi tridactyl, the median claw considerably longer, more slender, and erect than the

lateral claws.

TYPE SPECIES: *Lassenia lasseni* new species.


REMARKS: The forms described under this genus have many features in common, but there are also a few in which they differ markedly. It is possible that these may eventually necessitate the creation of a discrete genus for *L. spinifera*, although at present this does not seem necessary.

For a while it was felt that this genus and *Polydiscia* Methlagl 1928 might be identical. Judging from the original description of the latter genus, this might be so, but the possibility is sufficiently remote to make a final decision unwarranted until *Polydiscia squamata* has been fully described.


Lassenia lasseni new species

MALE: Body (Fig. 181) 1,534 to 1,716 μ long to tip of propodosoma, 936 to 1,144 μ wide, length/width = 1.44–1.64; average of five specimens 1,628 μ long, 1,066 μ wide, length/width = 1.53. Scutum (Fig. 183) broad, truncate posteriorly, rounded anteriorly, with a single pair of very slender smooth sensilla arising from coarse alveoli. Otherwise with seven to nine stiff, stout, setae on either side of the mid-line; these setae slightly fusiform and smooth. Crista metopica well developed, extending from posterior margin to a point a little more than half way between the alveoli of the sensilla and the anterior margin of the plate. Cuticle of scutum coarsely punctate except for the pale posterolateral portion of the plate. With the exception of the sensilla all of the setae of the scutum are of the same type. Ocular plates well developed and bearing two corneae, the anterior one the larger of the two. A single seta arising from a very minute sclerite between the ocular plate and the scutum; otherwise dorsum of propodosoma devoid of setae. Dorsal body setae (Fig. 184) stiff, smooth, straight, each arising from a small sclerite. Cuticle of dorsum smooth, without striae.

Coxa I (Fig. 192) with 45 to 50 smooth, slender setae plus the short, blunt, spikelike supra-



FIGS. 180–188. *Lassenia lasseni* n. sp., male: 180, gnathosoma, ventral; 181, dorsum; 182, venter; 183, propodosoma; 184, dorsal hysterosomal setae; 185, genital area; 186, legs I and II; 187, legs III and IV; 188, tip of rostrum, ventral.



FIGS. 189-201. *Lassenia laseni* n. sp.: 189, specialized setae of tarsus I, male; 190, tibia and tarsus of palp, male, posterior; 191, tibia and tarsus of palp, male, anterior; 192, coxae I and II, male; 193, coxae III and IV, male; 194, vestigial and other setae of tibia I, male; 195, *Lassenia*-organ of right and left sides of same male; 196, same, a different male; 197, intercoxal setae, male; 198, ventral hysterosomal setae; 199, *Lassenia*-organ, female; 200, Chelicera, male; 201, palp, male, anterior.

coxal seta; II with 40 to 45 setae. Dorsal portion of both coxae I and II membranous. Pars medialis coxae with 8 to 10 setae, outside of which are 20 to 25 setae in the intercoxal area. All of these plus a very few of the most medial setae of the coxae are two- to four-pronged, and quite stiff (Fig. 197). Apodemes behind II lying in a portion of the cuticle devoid of setae. Just anterior to III is the homologue of the unusual sclerite in the same position in the larva. It is extremely variable in form, but its constancy indicates that it is an important structure. In most cases it appears to be a plate with a gland opening. It is also present in the female. Coxae III and IV with about 40 setae each, all but one or two of which are slender, smooth and tapering; the remaining one or two are faintly barbed (Fig. 193). Cuticle of coxae brown in color, minutely and densely punctate. Genital sclerites (Fig. 185) forming a nearly hemispherical protuberance on the ventral surface of the body; deeply pigmented, each bearing about 24 smooth, slender setae. Paragenital sclerites slender, crescentic, each bearing 7 setae of the same form as those on the genital sclerites. Between the anterior ends of the paragenital sclerites is a hemispherical, deeply pigmented bulla. All sclerites associated with the genital opening minutely punctate. Three pairs of genital acetabula present; penis scarcely more than half the length of the genital sclerites. Anal sclerites well developed but slender, each bearing zero to five smooth setae. Most ventral setae much more slender and tapering than those on dorsum, each borne on a separate minute sclerite. Setae between coxae III of right and left sides of much the same form as those between coxae II, but not quite so heavy (Fig. 198).

Base of gnathosoma with 12 to 15 smooth


slender setae (Fig. 180), rostrum with 7 or 8 more in addition to those at the end of the rostrum. Supracoxal setae rather long (20 to 25 μ), tapering to a sharp point, inserted in the soft membranous dorsal cuticle of the coxae. Velum appearing elliptical in ventral view, inner margin fimbriated. Lateral to the velum is a pair of longer slender protorostral setae directed anteriorly, while just above the posterolateral margin of the velum is a pair of small spikelike deutorostral setae and a longer pair of tritorostral setae oriented in a posteroventral direction (Fig. 188).

Chelicerae (Fig. 200) relatively long and slender, heavily sclerotized, cuticle densely punctate; tarsus with 14 to 15 very minute teeth visible under high magnifications; chelicer al membrane pointed, extending to about middle of tarsus. Palpi (Fig. 201) relatively long, slender, straight. Trochanter devoid of setae, anterior aspect fenestrated. Femur with 29 to 30 setae, patella with 9 to 10 setae, anterior surface bare; tibia with about 10 normal setae plus the heavy spiniform terminal and subterminal setae. Tarsus (Figs. 190, 191) with an annulate solenidion at 0.4 to 0.5 *pd*, six to eight large eupathidia on distal half of segment and two to six normal setae (quite variable). All normal setae of palp appearing smooth at low magnification (100 x), but at higher magnifications a few very minute barbs can be seen on some of the setae. No tracheal openings could be seen on the dorsum of the gnathosoma in the position normally occupied by these, and no tracheae could be seen internally.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, n = normal setae, m = many).

Eupathidia present dorsally or marginally

	tf				pa				ti				ta					
	e	s ₃	e	v	s ₃	s ₄	e	v	s ₁	s ₄	s ₂	s ₃	e	f				
I	2	7	4	1	27	0?	5,7	1	(—m—)			0	m	1				
II	1	4	3	1	10	1	2	0	(-13-)	1	0	4	1					
III	1	5	2	0	10	0	1	0	0	8	0	0	3	0				
IV	1	6	2	0	8	0	1	0	0	10	0	0	3	0				



FIGS. 202-207. *Lassenia laszeni* n. sp., female: 202, solenidion₃, tibia I; 203, s₃, patella I; 204, s₁, tarsus I; 205, s₄, tarsus III; 206, s₄, tarsus II; 207, s₂, tarsus II.

FIGS. 208-214. *Lassenia laszeni* n. sp., larva: 208, dorsum; 209, Lassenia-organ; 210, palp, posterior; 211, chelicera and apodemes; 212, distal seta of tibia and basal seta of tarsus of palp; 213, palp, anterior; 214, propodosoma.

on all segments of all legs beyond the basifemur, none ventral, except on the tarsi. Vestigial setae present only on patella I and II and tibia I. Famulus of tarsus I at $0.55pd$, famulus₂ at 0.62 to 0.79 (this variation was noted on the right and left sides of a single male). Eupathidia of tarsus I extending from $0.17v$ to $0.17d$, claw fossa absent. Eupathidia of tarsus II extending from $0.43d$ to $0.97v$; tarsus tapering rapidly beyond $0.75d$, but no true claw fossa present. Tarsus III with eupathidia at $0.41d$, $0.64p$, and $0.96pv$; IV with eupathidia at $0.43d$, $0.48d$, and $0.67p$, none near end of tarsus. Solenidia of tarsi difficult to interpret because of convergence in form, especially between types s_1 , s_2 , and s_4 . No solenidia referable to type s_3 found on the tarsi. Tarsus II with a somewhat heavier solenidion, presumably s_2 , at $0.47d$, plus about 12 others presumably s_1 or s_4 or both. Claws of all tarsi simple, smooth, scythe-shaped.

FEMALE: Body 1,378 to 2,574 μ long, 806 to 1,872 μ wide, length/width = 1.38 to 1.71; average of six specimens 1,846 μ long, 1,235 μ wide, length/width = 1.50. Resembling the male in virtually all respects, even the genital sclerites and paragenital sclerites being only slightly larger and with possibly a very few more setae than in the male. Genital acetabula distinct, the anterior pair the largest. Anal sclerites also as in male.


LARVA: Body (Fig. 208) 522 to 576 μ long in partially engorged specimens. Scutum (Fig. 214) with anterior sensilla short, stiff and faintly pectinate; posterior sensilla very long, slender, smooth, uniformly tapered. Setae at anterolateral and posterolateral angles of scutum stiff, faintly roughened. Ocular plates with two distinct corneae. Cuticle of ocular plates and of scutum faintly punctate. Crista metopica completely absent. With 48 to 53 dorsal and marginal setae each borne on individual sclerites in the two specimens examined; lateral setae more slender than the others. Cuticle of dorsum without striae.

Coxae I and II with two setae each, III with three (occasionally four) simple smooth

setae; intercoxal area completely devoid of setae. Coxae I and II close together, urpore well developed; a structure of unknown nature (the "*Lassenia* organ," Fig. 209) laterally between II and III. Under oil immersion this appears to be a plate of much the same form as the setigerous sclerites, with the cuticle faintly punctate, and with an alveoluslike structure on the inner half of the plate. Near the center of this alveoluslike structure is a minute pore about 0.9 μ in diameter, which opens into a delicate chitinous duct about 22 μ long. At its inner end this duct expands to form a delicate cup or sphere approximately 3.6 μ in diameter. The duct is not hydrolyzed by the enzymes used to clear the mites.

Supracoxal setae present in the membranous dorsal portion of coxa I, but absent on II and III. Postcoxal area with 23 to 27 setae exclusive of those borne on the anal sclerites, all of these setae arising from individual feebly developed sclerites with faintly punctate cuticle (Fig. 221). Anal sclerites moderately developed, with two to three pairs of setae.

Base of gnathosoma (Fig. 229) devoid of setae, cuticle densely and minutely punctate, posterior half partly divided by vertical chitinous septum. Protorostral and tritorostral setae long and slender, deutorostral setae short and spikelike, posterorostral setae absent. Supracoxal setae rather long and uniformly tapered. Velum very delicate, semi-circular in form, consisting of a fimbria of extremely delicate chitinous processes extending inward toward the mid-line at the tip of the rostrum. Chelicerae (Fig. 211) with digitus mobilis a soft membranous process, tarsus with distal portion slender, scythe-shaped, of fairly uniform thickness throughout most of length, and typically with two small teeth near the end of the tarsus. Trochanter of palp (Figs. 210, 213) very short, ringlike, lacking setae. Femur greatly swollen and bearing five smooth setae on dorsal and posterior aspects. Patella incompletely separated from femur posteriorly (Fig. 210); with a single dorsal



FIGS. 215-217. *Lassenia laseni* n. sp., larva: 215, tarsus I, posterior; 216, tarsus II, posterior; 217, tarsus III, posterior.

seta. Distal portion of tibia curved sharply toward median line (Fig. 214); three pairs of smooth normal setae plus the greatly enlarged bifid distal seta. The latter is marked with somewhat irregular longitudinal striae (oil immersion, Fig. 212) and the two teeth are hollow in the distal half. Tarsus with two strongly curved, usually barbed heavy setae at base, a solenidion at 0.25 to 0.33*p*, one terminal and one subterminal eupathid; otherwise with five smooth, normal setae. Podocephalic canals reaching from cheliceral apodemes to, or nearly to, the urpore; a number of branches present.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathida, f = famulus, v = vestigial setae, c = companion setae, n = normal setae).

Femora I to III undivided (Figs. 218–220). Patella I and II with a vestigial seta dorsally. Tibiae with vestigial seta present only on I but with numerous solenidia dorsally as shown in the accompanying table; normal setae also abundant. Solenidion₄ on tibia I is longer, larger, and different in structure from the solenidia₃; moreover it has a companion seta at the base. Tarsus I with s₁ at 0.08 to 0.10*d* and a large eupathid at 0.64 to 0.69*d*, each with a companion seta; famulus at 0.24 to 0.27*pd* and a second eupathid at 0.91*pd*. The famulus is borne on a vesicular alveolus and is flattened and expanded at the tip (Fig. 228). Tarsus II with a delicately annulate solenidion at 0.34 and a eupathid with companion seta at 0.65*d*. Famulus like that on I but at 0.49*pd*; a second eupathid at 0.90*pd*. Tarsus III with a single eupathid at 0.91*pd* but with no other specialized setae. Most of the normal setae of the legs are delicately pectinate. All tarsi with three claws, the median one longer and more slender than the lateral ones.

TYPE LOCALITY: Mount Lassen, California, in a small cascade stream about one mile above Hat Lake; elevation about 7,250 feet (holotype male). July 9, 1954, and August 7, 1955. I. M. Newell, collector. Also collected at headwaters of Kings Creek, Mt. Lassen, California, August 6, 1955, by the writer.

REMARKS: This species can be differentiated from *L. scutellata*, new species, by the much larger number of setigerous sclerites on the dorsum (*L. scutellata*: only about 20 dorsal and marginal setae), the bifid form of the terminal seta of the palpal tibia (*L.s.*: this seta not bifid), and by numerous points of difference in the chaetotaxy of the legs which will be apparent by a comparison of the tables and descriptions given for the two species.


Lassenia lasseni is definitely a cold-stenothermal species and its distribution is undoubtedly limited by this fact. The larvae are parasitic upon small species of Diptera resembling Drosophilidae, found breeding along the banks of mountain streams. They are of the self-detaching type.

The correlation between the larva and adult of this species has been established solely on the basis of their occurrence at the same locality. While the correlation is to a certain degree provisional, it is reasonably certain that the two forms are conspecific. In all, 9 larvae, 6 nymphs, and 25 adults were collected at this location, and there is no indication that more than one species is involved in the collection. The species is not as common as the above numbers might indicate, for this represents the total catch of about 15 hours of hand collecting over a three-day interval. The adult was first collected July 9, 1954, and larvae and adults were found August 7, 1955.

Lassenia spinifera new species

FEMALE: Two specimens available for study

	tr	fe	pa			ti					ta					
	n	n	s ₃	v	n	s ₃	s ₄	v	c	n	s ₁	s ₂	e	f	c	n
I	1	10	2,3	1	8	12	1	1	1	16	1	0	2	1	2	50
II	3	10-12	0	1	8	4,5	0	0	0	17	0	1	2	1	1	43
III	3,4	9,10	0	0	8	3	0	0	0	15	0	0	1	0	0	40



FIGS. 218-229. *Lassenia lasseni* n. sp., larva: 218, leg I, trochanter to tibia; 219, leg II, trochanter to tibia; 220, leg III, trochanter to tibia; 221, venter; 222, solenidion₃, tibia I; 223, s₃, patella I; 224, s₃, tibia III; 225, s₄, tibia I; 226, s₂, tarsus II; 227, s₁, tarsus I; 228, famulus₁, tarsus I; 229, gnathosoma.

measured 1,326 and 1,794 μ to tip of scutum, 637 μ to 832 μ wide (length/width 2.08, 2.16). The smaller specimen was evidently an immature female and even the larger one was

nonovigerous. Scutum (Fig. 232) much more elongate than in the type species of the genus, and with a distinct anterior spine (hence the name). A pair of very elongate, slender,

smooth sensilla at or slightly behind the middle of the plate and a pair of much shorter setae near the base of the spine. These are of the same general form as the other setae of the dorsum of the propodosoma, but their position and orientation indicate that they are the anterior sensilla. Their alveoli are indistinct and very small in contrast with those of the other dorsal propodosomal setae which are well formed. Limit of scutum indistinct in anterolateral portions where it is all but impossible to trace in undissected specimens. Crista metopica well developed up to a point about half way between the posterior and anterior pairs of sensilla. Ocular plates elongate, bearing two pairs of prominent corneae, the posterior of which are the larger. In addition to the sensilla, the dorsum of the propodosoma bears 23 to 37 smooth, slender tapering setae on each side, 8 to 15 of which are definitely outside the scutum and 13 to 20 are definitely inside the scutum; one to three setae near the anterolateral margins of the scutum of uncertain position. Dorsal hysterosomal setae markedly different from those of the propodosoma, resembling those of *Lassenia lasseni*, stiff, rodlike, each borne on a small sclerite visible under high magnification (Fig. 246). The sclerites are not noticeably elevated. Membranous cuticle of hysterosoma completely smooth.

Coxae I with 65 to 75 smooth setae each (Fig. 234) plus the short peglike supracoxal seta in the membranous portion of the dorsal surface of the coxa. Coxa II with 35 to 50 setae; dorsal wall of distal part of coxa not sclerotized, but membranous like I. Both coxae I and II devoid of setae on their medial aspects. Intercoxal area with about 70 to 75 long stiff setae arranged in a rather characteristic radiating pattern; pars medialis coxae completely absent. Behind the sclerotized part of coxa II is a single to triple row of setae like those on the coxae, behind which is the usual transverse band of membranous cuticle devoid of setae. The presence of setae in the membranous area directly behind coxa

II, and the absence of the pars medialis coxae indicate a considerable reduction in the degree of sclerotization of the coxae of this species. While the specimen studied was not a completely matured female, it seems probable that the conditions described above would be found in ovigerous females also. Coxae III with 30 to 36 setae, IV with 65 to 75 setae (Fig. 244). Genital sclerites moderately well developed but not of the strongly crescentic form found in *Lassenia lasseni*, bearing 21 or 22 setae in a single to double row (Fig. 233). Paragenital sclerites very feebly developed, scarcely visible at low magnification, but at higher magnifications apparent as a slight change in texture of the cuticle; each sclerite bearing 12 to 15 setae. Anterior to the genital opening is the characteristic bulliform structure found also in the type species of the genus, and just anterior to this is a very small, transverse sclerite. Three pairs of genital acetabula present. The ventral body setae surrounding the genital opening are largely oriented in the direction of the genital opening, while the setae on the genital and paragenital sclerites are mostly oriented in a more ventrad direction. Anus (Fig. 240) completely terminal, anal sclerites fairly well delimited, weakly crescentic, bearing 16 to 17 setae each. Ventral hysterosomal setae borne on minute sclerites; shaft of seta considerably more slender and tapering than in the case of the dorsal setae, the two types intergrading laterally. Ventral cuticle mostly smooth, with only scattered traces of striations. Two small apodemes in the transverse strip of cuticle behind each coxa II. No trace of the characteristic gland found anterior to coxae III in the adults of *Lassenia lasseni* was seen in this species, but the material was not entirely favorable for study of this feature.

Base of gnathosoma (Fig. 248) totally devoid of setae, except for the supracoxals. Rostrum with four pairs of setae as shown in Figure 235. Deutorostrual setae short and peglike, concealed behind the margin of the velum in ventral view. The lateral arms of the

gnathosoma are quite elongate. Chelicerae (Figs. 245, 247) with posterior end bent rather sharply downward; tarsus with about four dorsal teeth in distal half, and four to five subdorsal teeth in basal half. Palpi (Fig. 242) considerably larger and relatively thicker than in *Lassenia lasseni* (Fig. 201). Anterior aspect of trochanter deeply dissected, but the sclerotization of the femur extends deep into this, so that the fenestration of the anterior wall of the trochanter is markedly reduced; left trochanter of holotype female with, right trochanter without, a seta on posterior aspect. Femur with 51, patella 36, and tibia with 27 smooth slender setae in the single specimen studied, plus the large terminal and smaller subterminal spiniform setae on the tibia. The terminal seta of the tibia is unidentate. Tarsus (Fig. 238) with a single solenidion at 0.88*pd*, this showing faint spiral structure and a somewhat swollen base. Proximal two-thirds of tarsus with 24 normal setae like those on the other segments of the palp; distal one-third of tarsus with 15 eupathidia showing distinct annulate structure internally. The eupathidia are somewhat heavier than the solenidion, and the curvature of the two types of setae is reversed.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, n = normal setae, m = many).

All solenidia of patella and tibia of all legs of *s*₃ type, except for one questionable *s*₄ on tibia I and II. Tarsus I with famulus at 0.80*pd* and many eupathidia extending from 0.13*v* to 0.17*d*; six or seven *s*₃ near the base of the tarsus, but all of the other many solenidia are more or less of the same form and size, so that it is impossible to differentiate the types with certainty. End of tarsus I rather abruptly


cut off, beginning at about 0.90*pd*; claw fossa absent. Tarsus II with famulus at 0.67*pd*, and about five eupathidia between 0.17*d* and 0.97*pv*; no ventral eupathidia. One seemingly distinct *s*₂ at 0.36*d*, one or two *s*₃ near the base of the tarsus, and two or three other solenidia of indeterminable type (*s*₁ or *s*₄). Tarsus II tapering rather abruptly from about 0.67*d* to the end of the segment, claw fossa absent. Tarsi III and IV each with about five *s*₃ or *s*₄; III with five eupathidia ranging from 0.19*d* to 0.95*v*, IV with an equal number ranging from 0.18*d* to 0.79*v*, terminal eupathid lacking. All tarsi with two heavy claws faintly hirsute along convex margin.

Telofemora well supplied with eupathids especially on I. Telofemur, patella, and tibia I notable in that the majority of the eupathidia are ventral in position. Of the 10 counted on telofemur I, 6 were dorsal or marginal in position, while 4 were ventral; of the 18 counted on patella I, 6 were dorsal or marginal while 12 were ventral; and of the 32 on tibia I only 8 were dorsal or marginal in position while 24 were ventral. These figures, like the rest of the figures in the accompanying table were based on counts made on a single specimen. Vestigial setae only on patella I and II and on tibia I; no vestigial seta on tibia II. All eupathidia on the telofemur, patella, and tibia of legs II to IV either dorsal or marginal in position, none ventral.

Lassenia scutellata new species

LARVA: Body (Fig. 249) 387 μ long, 243 μ wide, length/width 1.58 (one specimen only). Scutum subrectangular in form, divided into a small anterior portion bearing the anterior sensilla, and a much larger posterior portion bearing the posterior sensilla and other setae.

	tf		pa			ti				ta					
	e		s ₃	e	v	s ₃	s ₄	e	v	s ₂	s ₁	s ₄	s ₃	e	f
I	10		13	18	1	54	1?	32	1	(—m—)			7	±	1
II	3		7	4	1	8	1?	6	0	1	(-2,3-)		1,2	5	1
III	2		7	6	0	9	0	5	0	0	0	(-4,6-)		5	0
IV	1		8	8	0	17	0	6	0	0	0	(-4,6-)		6	0



FIGS. 230–238. *Lassenia spinifera* n. sp., female: 230, dorsum; 231, venter; 232, propodosoma; 233, genital area; 234, gnathosoma, coxae I and II, and intercoxal area; 235, tip of rostrum; 236, legs I and II; 237, legs III and IV; 238, tarsus of palp, posterior.

Cuticle of scutum slightly punctate; crista absent. Anterior sensilla only about half as long as posterior sensilla; scutum otherwise with two other pairs of setae. All setae of scutum with a few very minute barbs visible only under high magnification (400 x or higher). Ocular plates with two well-developed corneae, separated from the scutum by a narrow interval of striated cuticle. Dorsal and marginal setae numbering 20 pairs; venter with 7 pairs of ventral setae excluding those on the anal plates. Membranous cuticle of both dorsum and venter distinctly marked with striae which for the most part are parallel.

Coxae I (Fig. 251) each with two setae ventrally and a small inconspicuous supra-coxal seta; II with one, III with two setae. Between coxae I and II is a large well-developed urpore and anterior to coxa III at the margin of the body, a large sclerite obviously comparable with that found in *Lassenia lasseni*, new species. In the sole specimen available the soft parts were not very well hydrolyzed and a glandlike mass of cells can be seen extending up to the inner surface of the plate. Intercoxal area devoid of setae; postcoxal area with seven pairs of ventral setae surrounding the well-defined anal sclerites which bear an additional two pairs of slender setae.

Base of gnathosoma as described for *Lassenia lasseni*. Podocephalic canals probably present but not visible because of opacity of specimen. Supracoxal setae present, structure of velum obscured by organic accretions evidently resulting from feeding activities of mite; protorostral, deutorostral and tritorostral setae present, posterorostrals absent.


Chelicerae (Fig. 261) essentially as in *Lassenia lasseni*; tarsus with two teeth dorsally (not visible in figure because of rotation of

chelicera). Cuticle of chelicerae faintly and sparsely punctate, with parallel longitudinal striae dorsally. Palpi (Fig. 265) distinctly five-segmented, trochanter and femur completely separated, not partially fused as in *Lassenia lasseni*. Trochanter very short, ring-like, femur bearing only two setae, one dorsally and one anteriorly. Patella wedge-shaped, bearing a single long seta dorsally; tibia with only three very slender slightly pectinate setae in addition to the heavy terminal toothlike seta; the latter is not bifid at the tip but the microscopic structure of the hollow tip appears identical with that of the type species. Tarsus of palp with a distinctly annulate solenidion at 0.28*p*, one eupathid at 0.69*d*, and a second one terminally, otherwise with nine normal setae. The most basal of the latter is rather sharply deflexed and heavily pectinate, but it is not short and stout as in the genotype. Base of tarsus also with a pit-like structure dorsally, the exact nature of which is not known.

Chaetotaxy of legs approximately as shown in table (s = solenidia, e = eupathidia, f = famulus, v = vestigial setae, c = companion setae, n = normal setae).

Chaetotaxy of legs differing in numerous specific respects from *Lassenia lasseni*. Trochanters I to III (Figs. 262-264) with a single seta each. Femora with no trace of subdivision, so that the legs have only five free segments as in the type species. Patella I and II with vestigial seta of the same form as found on tibia I (Fig. 257). Tibia I with, II without a vestigial seta; tibiae not so richly supplied with solenidia₃ as in the case of *Lassenia lasseni*, there being only four on tibia I. Tibia I with a solenidion₄ at 0.72*d*, but II and III with only the solenidia₃. Tarsus I with s₁ and companion setae at 0.27*d*, a smooth, elongate spikelike famulus at 0.68*pd*, a eupa-

	tr	fe	pa					ti			ta					
	n	n	s ₃	v	n	s ₃	s ₄	v	c	n	s ₁	s ₂	e	f	c	n
I	1	6	2	1	5	4	1	1	1	8	1	0	2	1	2	36
II	1	6	1	1	4	2	0	0	0	9	0	1	2	1	1	29
III	1	5	1,0	0	4	1	0	0	0	8	0	0	1	0	0	24



FIGS. 239–248. *Lassenia spinifera* n. sp., female: 239, trochanter and femur of palp, anterior; 240, anus; 241 tibia and tarsus of palp, anterior; 242, palp, posterior; 243, tarsus I, showing setal types; 244, coxae III and IV. 245, chelicera; 246, dorsal hysterosomal setae; 247, end of chelicera; 248, gnathosoma, ventral.

thid and companion seta at $0.71ad$ and a second eupathid at $0.92p$ (Figs. 255, 258, 260). The famulus is inserted on a somewhat vesicular alveolus, but this is not as prominent as in *L. laseni*; moreover, the tip of the famulus tapers to a fairly sharp point and is not expanded at the tip as it is in the type species. Tarsus II with s_2 at $0.40d$, famulus at $0.61d$ of the same form as found on tarsus I,


a periepathid eupathid and companion seta at $0.67d$ and a second eupathid at $0.94p$. Tarsus III with a single eupathid at $0.92p$. Most normal setae of legs bihemipectinate, the pectinations visible at magnifications of 200 x and higher; total number of normal setae on legs about 145, compared with 240–245 in *L. laseni*. Other features of chaetotaxy as shown in the accompanying table. Three

claws on all tarsi, the median claw considerably more slender and erect than the lateral claws; lateral claws with a fringe of hairs along the outer margins, median claw also appearing hirsute under oil immersion.


TYPE LOCALITY: Spencer's Butte, Eugene, Oregon (holotype larva). The type and only specimen was found in material beaten from shrubs, July 10, 1952, collected by the writer. Since *Lassenia lasseni* has larvae of the self-detaching type, it is probable that the present species has this same habit. Therefore it is impossible to say whether the larva was actually crawling on the shrubs or whether it was

attached to a host insect which happened to be on the shrubs. It had obviously been feeding.

REMARKS: It is possible that this is the larva of *L. spinifera*, but since the locality records for the two are quite different, it is best not to correlate the larva and adult at this time. *L. scutellata* and *L. spinifera* are at present known only from the state of Oregon. *L. scutellata* can be readily differentiated from *L. lasseni* on the basis of the characters discussed under the remarks following the description of the type species.



FIGS. 249-254. *Lassenia scutellata* n. sp., larva: 249, dorsum; 250, gnathosoma; 251, venter; 252, tarsus of palp, posterior; 253, tarsus of palp, distidorsal; 254, cyst-like object of unknown nature attached to palpal eupathid, probably fungus or protozoan.



FIGS. 255–261. *Lassenia scutellata* n. sp., larva: 255, tarsus I, posterior; 256, famulus, tarsus I; 257, vestigial seta, tibia I; 258, tarsus II, posterior; 259, organism of unknown nature attached to seta at base of femur I; 260, tarsus III; 261, chelicera.

Polydiscia Methlagl 1928

The resemblance between the larva of *Polydiscia squamata* Methlagl 1928 and that of *Lassenia scutellata* n. sp. is so close as to leave little doubt of the very close relationship between *Polydiscia* and *Lassenia*. The genus must

therefore be removed to the Johnstonianidae. Thor and Willmann (1947, p. 342) placed the genus in the Trombiculinae, Radford (1950, p. 99) placed it in the Microtrombidiinae, and Baker and Wharton (1952, p. 251, *Polydiscia*, sic) listed it with the Trombidiidae whose subfamilial relationships are unknown.

The major points of similarity to *Lassenia* are the presence of two pairs of sensilla on the scutum (not a single pair as assumed by various authors), the large setigerous sclerites covering the dorsum, the 2-1-2 setal formula of the coxae, the presence of distinct setigerous anal sclerites, and the tridactyl tarsi of the larva. There are however, certain differences such as the apparently rectangular form of the base of the gnathosoma, the simple form of the palpi, the absence of a clawlike seta on the tibia of the palp and the subdivision of the femora of the legs, (provided these were all accurately represented in the original descriptions, which is improbable).

The fact that the anterior pair of setae of the scutum were not considered sensilla undoubtedly stems from the weak development of the alveoli of these setae. Precisely the same situation is found in *Lassenia scutellata* (Fig. 249), although close study of these setae at high magnification shows that they do have somewhat more elaborately developed alveoli than do the other setae of the scutum. The anterior setae of *Lassenia lasseni* are even less sensillar in form, although there is no doubt whatever that these are the true homologs of the anterior sensilla of such forms as *Diplothrombium*.

Thor and Willmann (1947, p. 343) stated that this was "wahrscheinlich ein Trombidioseerreger." Methlagl (p. 247) stated that *Polydiscia squamata* was responsible for trombidiasis in the vicinity of Vienna, although he earlier (p. 225) indicated that he had never found it on any host. Apparently the correlation between trombidiasis and *Polydiscia* was based solely on the seasonal abundance and general appearance of the latter. But this type of correlation can lead to absurd results as Methlagl himself (p. 247) so clearly showed when, on similar evidence, he supposed *Hauptmannia longicollis* Oudemans to be the larva of *Anystis vitis* (Schrank). For the present we can only conclude that *Polydiscia squamata* has not been shown to play any role in trombidiasis of man. On the contrary, its similarity to *Lassenia scutellata* (and hence to *L. lasseni*)

is good presumptive evidence that larvae of *P. squamata* are parasites of insects and not of vertebrates.


Crossothrombium Womersley 1939

This South Australian genus has been generally overlooked in recent catalogs (Thor and Willmann, 1947; Radford, 1950; Baker and Wharton, 1952). As Womersley pointed out, it is allied to *Johnstoniana*, and therefore belongs in the Johnstonianidae. However, contrary to Womersley's diagnosis, the holotype, *Crossothrombium parkhousei* Womersley 1939, appears to have two pairs of sensilla. The anterior pair, as in the case of *Lassenia*, is appreciably different from the posterior pair, but nevertheless they must be regarded as the homologs of the anterior sensilla. Womersley indicated that the eyes are absent which would be a significant point of difference between this genus and all of the others of Johnstonianidae. He also indicated that the dorsal setae arise from large "pits or circles," which are evidently the setigerous sclerites so typical of the Johnstonianidae. The tarsus of the palp was described as being "without terminal spines." If "spines" in this case means eupathidia, this would be unique in the family. This, too, requires verification. The description of both the genus and the species are too fragmentary to permit comparison with other genera of the family. One of the characters that can be seen in the figures is that there are several setigerous sclerites lying lateral to the scutum.

The species was described from a single specimen collected in South Australia, and was subsequently recorded from Victoria, Australia (Womersley 1942, p. 172).

Myrmicotrombium Womersley 1934

The original diagnosis of this genus was as follows: "Crista short, with two sensillary areas at anterior and posterior end. One eye on each side, in front of the anterior end of crista. Body as in *Microtrombidium*. Tarsi at end truncate, with one or two small, raised prom-



FIGS. 262–266. *Lassenia scutellata* n. sp., larva: 262, leg I, trochanter-tibia; 263, leg II, trochanter-tibia; 264, leg III, trochanter-tibia; 265, palp, posterior; 266, dorsal eupathid, famulus, and companion seta, tarsus I.

inences from which arise plain setae.” Womersley pointed out in his remarks that the presence of two pairs of sensilla on the scutum indicated a relationship to *Diplothrombium* and also to *Johnstoniana* (= *Robaultia*). However, a study of his figure of the scutum shows that this is in no way similar to the scutum of either *Diplothrombium* or of *Johnstoniana*. In fact it is very suggestive of the scutum of Erythraeidae such as *Balaustium*. Likewise the sharply cutoff form of the tarsus, with the distal face of the tarsus nearly vertical is not found in any Johnstonianidae known to the writer, but is found in some of the smaller Erythraeidae. Again, the setae are quite elaborately pectinate, which is also unlike the Johnstonianidae, in which the setae of the

body are generally smooth and borne on elevated alveoli. The single eye is also more characteristic of certain of the Erythraeidae than the Johnstonianidae. The association of the type species with ants would also be quite unusual for the Johnstonianidae which are largely subaquatic, and are normally found only in very wet situations which would be incompatible with ant nests. In view of these considerations it is the feeling of the writer that the genus *Myrmicotrombium* does not belong in the Johnstonianidae, but rather in the Erythraeidae. A study of the chelicerae of the type species would resolve this question; unfortunately Womersley did not describe these. The type species, *Myrmicotrombium brevicristatum* Womersley 1934 is known from a single

specimen collected at Glen Osmond, South Australia. The type is in the South Australian Museum.

REFERENCES

- BAKER, E. W., and G. W. WHARTON. 1952. *An Introduction to Acarology*. MacMillan Company. xiii + 465 pp.
- BERLESE, ANTONIO. 1887. *Acari, Myriapoda, et Scorpiones hucusque in Italia reperta*. Fascicle 42. Padova.
- . 1912. Trombidiidae. Prospetto di genere e delle specie finora noti. *Redia* 8: 1-291.
- . 1918. Centuria quarta di Acari nuovi. *Redia* 13: 115-192.
- COOREMAN, JEAN. 1952. Biologie et études faunistiques des Acariens. *Ninth Internatl. Cong. Ent., Trans.* 1: 1095-1100.
- GRANDJEAN, F. 1935. Les poils et les organes sensitifs portés par les pattes et le palpe chez les Oribates. *Soc. Zool. de France, Bul.* 60(1): 6-39.
- . 1947. Étude sur les Smarisidae et quelques autres Érythroïdes (Acariens). *Arch. de Zool. Expt. et Gén. Notes et Rev.* 85(1): 1-126.
- METHLAGL, A. 1928. Über die Trombidiöse in den österreichischen Alpenländern. *Akad. der Wiss. Wien, Math.-Nat. Kl., Denkschr.* 101: 213-249.
- MURRAY, ANDREW. 1876. Aptera. IN *Economic Entomology*. Scribner, Welford, and Armstrong. New York.
- OUDEMANS, A. C. 1937. *Kritisch Historisch Overzicht der Acarologie*. Gedeelte 3, Band D: Parasitengona Oudms. V. 1909. Pp. i-xix, 1349-1998. E. J. Brill. Leiden.
- RADFORD, C. D. 1950. *Systematic Check List of Mite Genera and Type Species*. 232 pages. Paris.
- THOR, SIG. 1935. Übersicht und Einteilung der familie Trombidiidae, W. E. Leach 1813 in Unterfamilien. *Zool. Anz.* 109(5/6): 107-113.
- THOR, S., and C. WILLMANN. 1947. Trombidiidae. IN *Das Tierreich, Lfg.* 71b: 187-541.
- WHARTON, G. W. 1948. Four new Peruvian Chiggers (Acarina, Trombiculidae), *Psyche* 55(2):87-100.
- WHARTON, G. W., D. W. JENKINS, J. M. BRENNAN, H. S. FULLER, G. M. KOHLS, and C. B. PHILIP. 1951. The terminology and classification of trombiculid mites (Acarina, Trombiculidae). *Jour. Parasitol.* 37(1): 13-31.
- WOMERSLEY, H. 1934. A revision of the trombid and erythraeid mites of Australia with descriptions of new genera and species. *Rec. So. Austral. Mus.* 5(2): 179-254.
- . 1939. Further notes on the Australian Trombidiidae with description of new species. *Roy. Soc. So. Austral., Trans.* 63(2): 149-166.
- . 1942. Additions to the Acarina of Australia (Trombidiidae and Calyptostomidae). *Rec. So. Austral. Mus.* 7(2): 169-181.